



**RAFFLES INSTITUTION**  
**YEAR 6 H2 CHEMISTRY 2016**  
**Term 3 Revision –**  
**Physical Chemistry Summary Notes**

---

## **Topics**

- (A) Stoichiometry & Gases
- (B) Atomic Structure
- (C) Chemical Bonding
- (D) Chemical Energetics
- (E) Reaction Kinetics
- (F) Chemical Equilibrium
- (G) Acid–Base Equilibria
- (H) Solubility Equilibria
- (I) Electrochemistry

## \* Quick Revision Notes \*

### A1. ATOMS, MOLECULES & STOICHIOMETRY and REDOX EQUATIONS

- 1) Know the definitions of the following terms!
  - (i) Relative Atomic Mass based on the  $^{12}\text{C}$  scale.
  - (ii) Relative Isotopic Mass based on the  $^{12}\text{C}$  scale.
  - (iii) Relative Molecular Mass based on the  $^{12}\text{C}$  scale.
  - (iv) Relative Formula Mass based on the  $^{12}\text{C}$  scale.
  - (v) Mole (in terms of Avogadro constant)
  - (vi) Empirical and Molecular formulae
- 2) Be able to calculate the relative atomic mass ( $A_r$ ) of an element given the relative abundances of its isotopes.

$$\text{Average } A_r = \frac{\text{abundance} \times A_r \text{ of isotope 1} + \text{abundance} \times A_r \text{ of isotope 2} + \dots}{\text{Total abundance}}$$

- 3) Calculate empirical and molecular formulae using combustion data or composition by mass.

For combustion of hydrocarbons:  $\text{C}_x\text{H}_y + (x + y/4)\text{O}_2 \longrightarrow x\text{CO}_2 + (y/2)\text{H}_2\text{O}$

Be careful when approximating after obtaining the ratio of moles.

E.g.

1.2	:	1	=	6 : 5
1.25	:	1	=	5 : 4
1.33	:	1	=	4 : 3
1.5	:	1	=	3 : 2
1.67	:	1	=	5 : 3
1.75	:	1	=	7 : 4

In general, multiply by suitable factor to get nearest whole number and DO NOT SIMPLY ROUND UP/DOWN.

Ending no.	.2	.25	.33	.5	.67	.75
Multiply by	5	4	3	2	3	4

- 4) Write and /or construct balanced equations and perform calculations using mole ratios.
 

Identify the limiting reagent!

1 mole of gas at r.t.p. and s.t.p. occupies 24 dm<sup>3</sup> and 22.4 dm<sup>3</sup> respectively.

No. of moles = mass/molar mass

Concentration in mol dm<sup>-3</sup> = no. of moles/volume of solution in dm<sup>3</sup>

Concentration in g dm<sup>-3</sup> = mass/volume of solution in dm<sup>3</sup>
- 5) 1 mole of a species contains  $6.02 \times 10^{23}$  particles of that species.
 

1 mole of NH<sub>3</sub> contains 1 mole of N atoms and 3 moles of H atoms,  $6.02 \times 10^{23}$  N atoms and  $3 \times 6.02 \times 10^{23}$  H atoms.
- 6) Be able to write balanced redox equations. (even if you've never seen the species before)
 

To write a balanced **half-equation**:

**Step 1:** Balance the **element** other than O and H.

**Step 2:** Balance **O** using H<sub>2</sub>O.

**Step 3:** Balance **H** using H<sup>+</sup>.

**Step 4:** Add equal no. of moles of OH<sup>-</sup> on both sides of half equation so as to completely neutralize all H<sup>+</sup>. (Step 4 is used only if in alkali/neutral conditions)

**Step 5:** Balance charges using electrons.

7) Be familiar with the following terminology:

Sulfur trioxide	SO <sub>3</sub>	Nitrogen dioxide	NO <sub>2</sub>
Sulfate/ sulfate (VI)	SO <sub>4</sub> <sup>2-</sup>	Nitrate/ nitrate (V)	NO <sub>3</sub> <sup>-</sup>
Sulfuric acid	H <sub>2</sub> SO <sub>4</sub>	Nitric acid	HNO <sub>3</sub>
Sulfur dioxide	SO <sub>2</sub>	Nitrogen oxide	NO
Sulfite/ sulfate (IV)	SO <sub>3</sub> <sup>2-</sup>	Nitrite/ nitrite (III)	NO <sub>2</sub> <sup>-</sup>
Sulfurous acid	H <sub>2</sub> SO <sub>3</sub>	Nitrous acid	HNO <sub>2</sub>
Chromate (VI)	CrO <sub>4</sub> <sup>2-</sup>	Dichromate (VI)	Cr <sub>2</sub> O <sub>7</sub> <sup>2-</sup>
Manganate (VII)	MnO <sub>4</sub> <sup>-</sup>	Manganese (II)	Mn <sup>2+</sup>

8) Be familiar with these various common titrations.

(a) MnO<sub>4</sub><sup>-</sup> with Fe<sup>2+</sup> (purple MnO<sub>4</sub><sup>-</sup> is reduced to colourless Mn<sup>2+</sup>, Fe<sup>2+</sup> is oxidized to Fe<sup>3+</sup>.)  
[MnO<sub>4</sub><sup>-</sup> : Fe<sup>2+</sup> = 1: 5]

(b) MnO<sub>4</sub><sup>-</sup> with H<sub>2</sub>O<sub>2</sub> (purple MnO<sub>4</sub><sup>-</sup> is reduced to colourless Mn<sup>2+</sup>, H<sub>2</sub>O<sub>2</sub> is oxidized to O<sub>2</sub>.)  
[MnO<sub>4</sub><sup>-</sup> : H<sub>2</sub>O<sub>2</sub> = 2: 5]

Important note: In this case the half equation for H<sub>2</sub>O<sub>2</sub> is H<sub>2</sub>O<sub>2</sub> → O<sub>2</sub> + 2H<sup>+</sup> + 2e<sup>-</sup> and NOT H<sub>2</sub>O<sub>2</sub> → H<sub>2</sub>O + O<sub>2</sub>, which shows the disproportionation of H<sub>2</sub>O<sub>2</sub> as it is oxidized to O<sub>2</sub> as well as reduced to H<sub>2</sub>O.

(c) Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> with Fe<sup>2+</sup> (orange Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> is reduced to green Cr<sup>3+</sup>, Fe<sup>2+</sup> is oxidized to Fe<sup>3+</sup>.)  
[Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> : Fe<sup>2+</sup> = 1:6]

(d) A common 2-step reaction involving iodine.

(i) KI (a source of I<sup>-</sup>), is oxidized by excess reagent to get I<sub>2</sub>

(ii) I<sub>2</sub> formed from (i) is then titrated with S<sub>2</sub>O<sub>3</sub><sup>2-</sup>, where I<sub>2</sub> is reduced to I<sup>-</sup> and S<sub>2</sub>O<sub>3</sub><sup>2-</sup> is oxidized to S<sub>4</sub>O<sub>6</sub><sup>2-</sup>. [I<sub>2</sub> : S<sub>2</sub>O<sub>3</sub><sup>2-</sup> = 1: 2]

(e) C<sub>2</sub>O<sub>4</sub><sup>2-</sup> as a reducing agent. How to write the equation for C<sub>2</sub>O<sub>4</sub><sup>2-</sup> being oxidized?  
Product: CO<sub>2</sub>. Use steps in point 7 above to write the half equation.

## A2. THE GASEOUS STATE

1) Be able to state basic assumptions of the kinetic theory of ideal gases, namely

i) Gases consist of small particles of negligible volumes.

ii) The gaseous particles exert no attractive forces on each other.

2) Know the conditions necessary for a gas to approach ideality (i.e. low pressure and high temperature). This is because

(a) at low pressure, the *volume of gas particles* of a real gas becomes negligible as compared to the volume occupied by the gas.

(b) at high temperature, the gas particles are able to overcome most of the intermolecular forces of attraction.


Note that at these conditions of low pressure and high temperature, the real gas is able to behave close to the assumptions of an ideal gas (in point 1 above). Conversely, at high pressures and low temperatures, a real gas will deviate most greatly from ideality.

3) Gases with larger molecules (and thus electron cloud size) and stronger intermolecular attractive forces tend to deviate more from ideal gas behaviour.

4) State & use the general gas equation pV = nRT in calculations, including the determination of M<sub>r</sub>.

Units are very important!!!!

Value of R	8.31 J K <sup>-1</sup> mol <sup>-1</sup>	0.082 atm <sup>-1</sup> mol <sup>-1</sup> K <sup>-1</sup>
Pressure (p)	Pa or Nm <sup>-2</sup> (1 atm = 1.013 x 10 <sup>5</sup> Pa)	atm
Volume (V)	m <sup>3</sup> (1 cm <sup>3</sup> = 1 x 10 <sup>-6</sup> m <sup>3</sup> ; 1 dm <sup>3</sup> = 1 x 10 <sup>-3</sup> m <sup>3</sup> )	dm <sup>3</sup>
Temperature (T)	K (25° = 25 + 273 K)	K



## C. CHEMICAL BONDING

### 1) When drawing dot-cross diagrams, note that

- All atoms in a molecule "want" to fulfil the OCTET rule – but this is NOT compulsory. E.g.  $\text{BF}_3$  has only 6 electrons in the valence shell of B, and that is why it readily accepts a lone pair from  $\text{NH}_3$  to form  $\text{BF}_3\cdot\text{NH}_3$  via coordinate (dative) bonding.
- General steps when drawing dot-cross diagrams for simple covalent molecules:
  - (i) Satisfy all side atoms first. E.g. Cl is a side atom and needs 1 more electron to be shared with it. Hence the central atom forms a single bond with it.
  - (ii) Account for all the electrons of the central atom. e.g.  $\text{XeF}_4$ : after forming 4 single bonds with F, I must draw 4 more electrons for Xe since it is from group 8 and should have 8 electrons around its valence shell.
  - (iii) Check that, if the element is in period 1 or 2, that the central atom does not have more than a total of 8 shared electrons around it.
- Elements in period 3 (period containing P) and beyond can expand their octet and thus accommodate more than 8 valence electrons. When it is possible to expand its octet, the central atom would form double bonds with a side atom that needs to share 2 electrons (e.g. O) rather than coordinate (dative) bonds even if it has to accommodate more than 8 valence electrons. This is because double bonds are stronger than a single dative bond.
- For compounds  $\text{XO}_n$ , draw all the X-O bonds as double bonds if X is from period 3 and beyond. Otherwise, dative bonds may be drawn. For anions  $\text{XO}_n^{m-}$ , each of the negative charge is accommodated on m number of oxygen atoms, and these are singly bonded to the central atom. n-m number of oxygen atoms are then drawn as double bonds (if X is from period 3 and beyond), otherwise dative bonds may be drawn.
- For ionic bonding, the correct number of each ion should be indicated.

Remember to indicate the overall charge of any ions outside the square bracket.

### 2) Predicting shapes of molecules/ions.

Be clear about the VSEPR Theory- Regions of electron density arrange themselves as far apart as possible so as to minimize repulsion.

Find out the electron pair geometry based on total number of electron regions, and then determine the shape based on number of bond pairs and lone pairs. Note: the shape of the molecule and how the bond and lone pairs are arranged determine whether a molecule is polar. Symmetrically arranged  $\Rightarrow$  zero overall dipole moment  $\Rightarrow$  non-polar

Factors affecting bond angles:

i)  $\text{l.p.-l.p. repulsion} > \text{l.p.-b.p. repulsion} > \text{b.p.-b.p. repulsion}$

Lone pairs exert greater repulsion as they are closer to the nuclei. E.g. 4 bond pairs: bond angle  $109.5^\circ$ ; 3 bond pairs 1 lone pair: bond angle  $107^\circ$ ; 2 bond pairs 2 lone pairs: bond angle  $105^\circ$

ii) Electronegativity

Atoms which are more electronegative than the central atom will draw the shared electron density further from the nucleus, resulting in a weaker repulsive force exerted by the bonding pair. Bond angle will be smaller.

iii) Size of the attached atoms

A larger attached atom will have a larger electron cloud which will exert a repulsive force and prevent the attached atoms from coming too close together. Bond angle will be larger.

3) When explaining m.p. or b.p.

Structure	Giant Metallic	Giant Ionic	Giant Covalent	Simple Covalent (weak forces of interaction between molecules although they have <b>very strong covalent bonds within the molecule</b> . It is the intermolecular interactions that are being overcome during melting /boiling, not the covalent bonds)		
Type of bonds being overcome when melting and/or boiling	Strong forces of attraction between positive metallic cations and sea of delocalised electrons	Strong ionic bonds between oppositely charged ions	Strong covalent bonds between all atoms	Instantaneous dipole-induced dipole interactions (id-id) - for non-polar molecules, and essentially present in all molecules	Permanent dipole-permanent dipole interactions (pd-pd) - for polar molecules	Hydrogen bonds - molecules with H <u>DIRECTLY</u> bonded to F/O/N.
Remarks	When comparing m.p. of metals, the smaller the cation and the larger the number of delocalised electrons, the stronger the bond. i.e. strength of metallic bond $\propto q^+/r^+$	Bond strength determined by lattice energy.	Bond strength dependent on effectiveness of overlap. e.g. melting pt of C (diamond) higher than m.p. of Si.	NOTE: Hydrogen bonds $>$ pd-pd $>$ id-id <u>ONLY</u> if the molecules have similar electron cloud sizes. E.g. compounds with id-id interactions will have higher m.p./b.p. than compounds with pd-pd interactions if the electron cloud size of the molecule with id-id interactions is larger than that of the molecule with pd-pd interactions.		
m.p	Generally high			Generally low		

4) Ionic bond –

- a) The strength of the ionic bond is indicated by the lattice energy of the ionic compound:

$$|L.E.| \propto |(q_+ q_-) / (r_+ + r_-)|$$

From the relationship above, we can deduce that high charges and small inter-ionic distances will lead to a **more exothermic** L.E. and hence, a stronger ionic bond.

- b) Ionic bonds are non-directional, i.e. there is no preferred orientation of the ions.
- c) Ionic compounds usually:
- Possess high m.p. and b.p. because all the ionic bonds need to be weakened / broken during a change of state.
  - Are soluble in a polar solvent (e.g. water) but insoluble in a non-polar solvent (e.g. hexane). For the ionic compound to be soluble, the energy **released** from solvation (forming ion-solvent bonds) **must** be comparable with the energy **absorbed** for breaking up the lattice ( $\propto$  L.E.).
  - Conduct electricity in the **molten** or **aqueous** states.

5) The covalent bond –

Usually formed between non-metallic atoms. Some **common exceptions**:  $\text{FeCl}_3$ ,  $\text{AlCl}_3$ ,  $\text{BeCl}_2$  are **covalent** even though formed between metallic and non-metallic atoms. This covalent character is a result of the high charge density of the cation and polarisability of the anion. ( $\text{AlF}_3$  is ionic due to the small anion which is not easily polarised).

The strength of the covalent bond is dependent on its bond energy (the energy required to break one mole of a particular bond in one mole of a gaseous substance) –

- The higher the bond order, the stronger the covalent bond, e.g.  $\text{C}=\text{C} > \text{C}-\text{C}$ .
- The **smaller the orbitals involved, the less diffused the orbitals and the more effective the overlap and the stronger the covalent bond.** (most significant reason to account for strength of covalent bonds)
- The greater the difference in the electronegativity between the bonding atoms, the more polar the bond, the stronger the covalent bond (in general) because the resultant dipoles may reinforce the bond through added electrostatic attraction between the atoms.

6) Be able to describe and draw  $\sigma$ - and  $\pi$ -bonds.

$\sigma$ -bonds are a result of head-on overlap,  $\pi$ -bonds are a result of side-on overlap. A single bond is made up of 1  $\sigma$ -bond, while a double bond is made up of 1  $\sigma$ -bond and 1  $\pi$ -bond and a triple bond is made up of 1  $\sigma$ -bond and 2  $\pi$ -bonds.

7) Possible ways of forming a dimer

- through dative (coordinate bonding) e.g.  $\text{Al}_2\text{Cl}_6$
- through hydrogen bonds e.g.  $(\text{HCOOH})_2$

8) Be able to account for the lower density of ice compared to water w.r.t. hydrogen bonding.

## D. CHEMICAL ENERGETICS

You have to know ALL the definitions VERY WELL!!!!

- ☐ Standard Enthalpy Change of *Reaction*,  $\Delta H_r^\circ$ :

The enthalpy change when molar quantities of reactants as specified by the chemical equation react to form products at 1 atmosphere pressure and 298 K temperature.

- ☐ Standard Enthalpy Change of *Formation*,  $\Delta H_f^\circ$ :

The enthalpy change when 1 mole of a pure compound in a specified state is formed from its constituent elements in their standard states, at 298 K and 1 atmosphere pressure.

- ☐ Standard Enthalpy Change of *Combustion*,  $\Delta H_c^\circ$ :

The heat energy *evolved* (ie exothermic) when 1 mole of the substance is completely burned in oxygen at 298 K and 1 atmosphere pressure.

- ☐ Standard Enthalpy Change of *Hydration*,  $\Delta H_{hyd}^\circ$ :

The heat energy *evolved* when 1 mole of the gaseous ion is dissolved in a large amount of water at 298 K and 1 atm.

- ☐ Standard Enthalpy Change of *Solution*,  $\Delta H_{soln}^\circ$ :

The heat change when 1 mole of solute is completely dissolved in enough solvent so that no further heat change takes place on adding more solvent (infinite dilution) at 298 K and 1 atm.

- ☐ Standard Enthalpy Change of *Neutralisation*,  $\Delta H_{neut}^\circ$ :

The heat change when an amount of acid neutralises a base to form 1 mole of water (in dilute aqueous solution) at 298 K and 1 atmosphere pressure.

- ☐ Standard Enthalpy Change of *Atomisation*,  $\Delta H_{at}^\circ$ :

The enthalpy change when 1 mole of atoms in the gas phase is formed from the element in the defined physical state under standard conditions.

- ☐ *Bond energy*

The **bond energy** of a X-Y bond is the average energy absorbed when 1 mole of X-Y bonds are broken in gaseous state. The stronger the bond, the higher the bond energy.

- ☐ *Bond dissociation energy*:

**Bond Dissociation Energy** of a X-Y bond is the energy required to break 1 mole of that particular X-Y bond in a particular compound in gaseous state.

- ☐ *Lattice Energy*

**Lattice energy** is the enthalpy change when one mole of a pure solid ionic compound is formed from its constituent gaseous ions under standard conditions.



- ☐ *First Ionisation Energy:*

**First ionisation energy** is the energy required to remove 1 mole of electrons from one mole of gaseous atoms in the ground state to form one mole of gaseous unipositive charged cations.

- ☐ *First Electron Affinity:*

**First electron affinity** is the energy change when 1 mole of electrons are added to one mole of atoms in the gaseous state to form one mole of gaseous  $X^-$  ions.

- ☐ *Second Electron Affinity:*

**Second electron affinity** is the energy change when 1 mole of electrons are added to one mole of gaseous  $X^-$  ions to form one mole of gaseous  $X^{2-}$  ions.

Summary of problem solving approach for Chemical Energetics

1.  $q = mc\Delta T$
2. Shortcuts: Use of formulae/ bond energies.
3. Hess' Cycle
4. Energy Level Diagram

1. **Step 1:** Using  $q = mc\Delta T$  to find heat energy released/ absorbed.

It is easy to identify when to use this equation. The question will mention that there is a rise in temperature in the solution, and immediately this equation should come to mind. However, many students do not understand what the symbols represent.

**(c)-Specific heat capacity of water**  
quantity of heat required to raise the temperature of 1g of water by 1 K. The heat energy released or absorbed is used to raise or lower the temperature of water. The mass is thus **mass of SOLUTION ONLY**.

**(m) mass**  
the mass that is computed must be the **mass of SOLUTION ONLY** (aqueous solutions consist mainly of water) and does not include mass of any solids.

**Step 2:** Finding enthalpy change of reaction using  $\Delta H = q/n$

Note:  $n$  = no. of moles of limiting reagent

2. Use of formulae/ bond energies  
(short-cut; actually can be derived using Hess' cycle).


- (a)  $\Delta H_{rxn} = \Sigma \Delta H_f(\text{products}) - \Sigma \Delta H_f(\text{reactants})$

Qn: When do we use this equation?

Ans: When data given is in the form of  $\Delta H_f$ .

Example: *(Pls verify your answers with your friends.)*

The yellow gas chlorine dioxide,  $\text{ClO}_2$ , has been used for many years as a flour-improving agent in bread-making. It can be made in the laboratory by the following reaction:



Calculate and comment on, the value of  $\Delta H_f^\ominus(\text{ClO}_2)$ , given the following data:

$$\Delta H_f^\ominus(\text{AgClO}_3) = -25 \text{ kJ mol}^{-1}$$

$$\Delta H_f^\ominus(\text{AgCl}) = -127 \text{ kJ mol}^{-1}$$

[3] N92/I/4(a)

**Important Note:**  $\Delta H_f$  of elements in their standard states = 0

(b) Using bond energy data from *Data Booklet*.

Qn: When do we use this method?

Ans: When the question asks you to use bond energy values from the *Data Booklet*.

Step 1: Draw out the FULL structural formulae of all species in the chemical equation. This is so that you can see all the bonds and not miss out on any bonds.


Step 2: Refer to *Data Booklet* for bond energy values. Write them in a table form.

Bonds broken (+ve)	Bonds formed (-ve)

Step 3: Remember that energy is required to break bonds and energy is released when bonds are formed. Therefore all values in the bonds broken column should be **positive**, while all values in the bonds formed column should be **negative**. To find the overall enthalpy for reaction in question, simply add the summation of the 2 columns together.

Example: (*Pls verify your answers with your friends.*)

Hydrogen is used in large quantities in industry to convert nitrogen into ammonia, for use in fertilizers. One method of manufacturing hydrogen is to pass methane and steam over a heated nickel catalyst.



- Use the value of  $\Delta H_1$  above, and bond energy values from the *Data Booklet*, to calculate the total bond energy in the carbon monoxide molecule.
- Suggest why the bond energy you have calculated in (i) is larger than either of the carbon-oxygen bond energies in the *Data Booklet*.

J96/I/2


Note: A very common question asked is why bond energy value differs from that found in the *Data Booklet*. Reason: Bond energy values in the *Data Booklet* are average values.

### 3. Hess' Cycle

Qn: When do we use this method?

Ans: For almost ALL thermochemistry questions except for those which specify that an energy level diagram must be drawn.

A Hess' cycle looks like this:



Step 1: Write down the equation for the enthalpy change in question. Reactants at A, products at B.

Step 2: **Decide what should be written for C.** You will decide what C should be depending on the **information that is provided**. If the question provides you with enthalpy of atomization values, for example, there will be atoms at C. If the question gives you enthalpy of combustion values, then you will have the products of the combustion of the species at A and B at C. Species at A, B and C must **all balance** with each other.

Step 3: Draw the arrows. The direction of the arrows will depend on the information given. This requires a thorough knowledge of the different enthalpy definitions. Plug in the values of the different enthalpies next to the arrows.

Step 4: Use your knowledge of vectors to work out the enthalpy change in question.

#### 4. Energy level diagrams

Qn: When do we use this method?

Ans: When the question asks explicitly for an energy level diagram.


Typically used when lattice energy is involved.

Thorough knowledge of enthalpy definitions needed.

Relevant data that can be obtained from Data Booklet:

- 1<sup>st</sup> ionization energy
- 2<sup>nd</sup> ionization energy
- bond energy

An energy level diagram looks like this, with varying number of steps/ energy levels:



The cycle must be complete. i.e. forms one full circle.

- Each step represents one enthalpy change.
- An arrow pointing upwards represents an endothermic reaction.
- An arrow pointing downwards represents an exothermic reaction.
- Labelling of numbers on y-axis not needed, just label each enthalpy change. If there is a 0 labelled on the y-axis, that would represent the **ELEMENTS at standard conditions**.

## E. Kinetics Summary

### 1. Know the definitions!

(i)	rate of reaction	The <b>change in concentration</b> of a reactant or product with time. i.e. rate at a given time $t = -\frac{d[\text{reactant}]}{dt} = \frac{d[\text{product}]}{dt}$ .
(ii)	rate equation	The rate equation or rate law is a mathematical equation that shows how the rate of reaction is dependent on the concentrations of the reactants; it relates the rate of the reaction to the concentration of reactants raised to the appropriate power. E.g. $\text{rate} = k[A]^m[B]^n$
(iii)	order of reaction	① The order of reaction with respect to a reactant is the power to which the concentration of that reactant is raised to in the rate law. ② The overall order of reaction is the sum of the powers to which the concentrations of the reactants are raised to in the rate law.
(iv)	rate constant	The rate constant, $k$ is a <b>proportionality constant</b> in the rate equation of the reaction.
(v)	half-life of a reaction	The time required for the concentration of a limiting reactant to decrease to half of its initial concentration.
(vi)	rate-determining step	The <b>slowest</b> step in the sequence of steps leading to the formation of the product.
(vii)	activation energy, $E_a$	The minimum amount of energy that the reactant particles must possess before they can collide successfully to form products
(viii)	catalyst	A substance which <b>increases the rate</b> of a chemical reaction without itself undergoing any permanent chemical change. It is <b>regenerated</b> at the end of the reaction.

### 2. Determining orders of reaction and rate equations

#### a. Observing initial rates data (Direct comparison)

If rate remains constant when concentration is doubled  $\Rightarrow$  zero order

If rate doubles when concentration is doubled  $\Rightarrow$  1<sup>st</sup> order

If rate quadruples when concentration is doubled  $\Rightarrow$  2<sup>nd</sup> order

(e.g. when concentration of a reactant is doubled while keeping concentration of the other reactants constant, rate doubles  $\Rightarrow$  order 1 with respect to the reactant)



#### b. Mathematical equations (Indices method)

Usually used because (a) is not feasible due to more complicated numbers

e.g. 
$$\frac{\text{initial rate of experiment 2}}{\text{initial rate of experiment 1}} = \frac{k[A]_2^a[B]_2^b}{k[A]_1^a[B]_1^b}$$

### c. Graphical data

Example:



Expt	$[\text{H}_2\text{O}_2] / \text{mol dm}^{-3}$	Initial $[\text{OH}^-] / \text{mol dm}^{-3}$
1	0.1	0.1
2	0.1	0.05

#### Solution

- To find order of reaction with respect to  $\text{H}_2\text{O}_2$ , use one of the graphs and find the half life. If half life is constant, order is 1 wrt  $\text{H}_2\text{O}_2$ .
- To find order of reaction with respect to  $\text{OH}^-$ , draw a tangent at  $t = 0$  for both expts and compare the value of the gradient (which represents rate) when  $[\text{OH}^-]$  in expt 2 is half that in expt 1. If grad of curve from expt 2 is half that in expt 1, order wrt  $\text{OH}^-$  is 1.

### 3. Pseudo-order reactions.

Consider a reaction in which  $\text{rate} = k[\text{A}][\text{B}]$

If  $[\text{A}]$  is essentially constant, then  $\text{rate} = k'[\text{B}]$  where  $k' = k[\text{A}]$ .

- This occurs
  - when **A** is present in large excess such that its concentration hardly changes
  - if **A** is a catalyst.
- This is a pseudo first order reaction wrt to **A**. If a graph of  $[\text{B}]$  against time was plotted,
  - the graph will have a constant half life ( $t_{1/2}$ )
  - $t_{1/2} = \ln 2 / k' = \ln 2 / (k[\text{A}])$ . The true rate constant  $k$  can be found.

### 4. Using rate law to determine mechanism

Verify that a suggested reaction mechanism is consistent with the experimentally observed kinetics. The slow step is the rate-determining step and must contain the correct ratio of reactants in the rate equation.

When predicting the order of reaction from a reaction mechanism, the rate equation should only contain reactants and NOT INTERMEDIATES.

### 5. Collision theory

Using microscopic processes to explain the phenomenon of reaction and its rate. Reactions only occur if there are effective collisions, i.e.

- reactant molecules colliding with  $E > E_a$  and
- the correct orientation.

Factors affecting the rate of a reaction:

<b>Concentration / Pressure</b>	: Higher concentration / pressure increases the frequency of effective collisions.
<b>Temperature (T)</b>	: At higher T, the fraction of reactant molecules with $E > E_a$ increases. Use Maxwell-Boltzmann Distribution curve to explain.
<b>Presence of catalyst</b>	: A catalyst provides an alternative reaction pathway of lower $E_a$ and more reactant molecules have energy greater than or equal to $E_a$ , leading to a higher frequency of effective collisions and hence a higher reaction rate. (use Maxwell-Boltzmann Distribution Curve to explain).

## 6. Catalysis

Two modes of catalysis:

- (a) Homogeneous (catalyst and reactants are in the same phase)
- (b) Heterogeneous (catalyst and reactants are in different phase)


### (a) Heterogeneous catalysis

#### (i) Important steps in heterogeneous catalysis

<b>Step 1</b>	<b><u>Adsorption</u></b> of the reactant particles onto the <b><u>active sites</u></b> on the surface of the catalyst. - Weak bonds form between the reactant particles and the catalyst. - Surface concentration of the reactants is increased.
<b>Step 2</b>	<b><u>Reaction</u></b> at the surface. - The <b>activation energy</b> of this process is lower than that of the uncatalysed reaction <b><u>because the bonds within the reactant particles are weakened by the adsorption effects</u></b> , thereby reducing the energy required to disrupt them - The reactant particles are brought into <b>close contact</b> and are <b><u>properly orientated</u></b> for reaction
<b>Step 3</b>	<b><u>Desorption</u></b> of the reactants or products from the surface.

#### (ii) Examples of heterogeneous catalysis


##### Haber Process




Conditions: 250 atm, 450°C, Fe

##### Catalytic converter

- As the gases enter, the oxides of nitrogen (NO and NO<sub>2</sub>) are reduced to N<sub>2</sub> by the excess CO present, with rhodium acting as the catalyst.





- The CO is also oxidised to CO<sub>2</sub> with platinum and palladium as the catalysts.



- The unburnt hydrocarbons are oxidised to CO<sub>2</sub> and H<sub>2</sub>O with platinum and palladium as the catalysts.

<b>(b) Homogeneous catalysis</b>
----------------------------------

- Reaction catalysed by  $\text{Fe}^{3+}$ :



The catalysed reaction proceeds via a 2-step route. Each step involves a reaction between oppositely charged ions which have a natural tendency to attract each other. This lowers the activation energy and enhances the reaction rate. ( $E_{\text{cell}} > 0$  for each step of the reaction)

- Enzymes catalyse reactions by providing an alternative reaction pathway with a lower activation energy. To bring this about, the enzyme forms a complex with the substrate or substrates (reactants) of the reaction. Thus a simple picture of the enzyme action is:



Once the products of the reaction are formed, they are released and the enzyme is free to form a new complex with more substrate.


## F. Chemical Equilibrium

### Reversible Reactions and Dynamic Equilibrium

1. Irreversible reactions are reactions that take place in one direction almost exclusively.

Eg: Burning of magnesium in air:  
 $2\text{Mg (s)} + \text{O}_2 \text{ (g)} \rightarrow 2\text{MgO (s)}$


Only the forward reaction takes place.



2. Reversible reactions are reactions that take place in both directions ( $\rightleftharpoons$ ) at comparable rates.

Eg: Rxn of  $\text{SO}_2$  with  $\text{O}_2$  :  
 $2\text{SO}_2 \text{ (g)} + \text{O}_2 \text{ (g)} \rightleftharpoons 2\text{SO}_3 \text{ (g)}$

Both the forward and backward reaction takes place simultaneously.



3. For reversible reactions, the system can reach a stage in which the **rate of forward reaction = rate of backward reaction** so that **no macroscopic changes are observed**. The system is said to be at equilibrium. Since microscopic changes are still occurring, the system is said to be in dynamic equilibrium.
4. Equilibrium is only achieved in a closed system. An open system which allows matter to escape or enter cannot reach equilibrium unless there is absolutely no gain or loss of materials from or to the surroundings.
5. Once equilibrium is reached, the system will remain in this state, until conditions affecting rate of forward/ backward reaction is altered. (eg, temperature, pressure, concentration of reactant/product)

### Law of Chemical Equilibrium and Equilibrium Constants

1. **Reaction Quotient** (or mass action expression)

(a) For a reversible reaction represented by  $a\text{A} + b\text{B} \rightleftharpoons c\text{C} + d\text{D}$ ,

$$\text{Reaction Quotient, } Q_c = \frac{[\text{C}]^c [\text{D}]^d}{[\text{A}]^a [\text{B}]^b}$$

[X] is instantaneous concentration (in  $\text{mol dm}^{-3}$ ) of X at time t

- (b) From time = 0 till time =  $t_{\text{eqm}}$ , the reaction quotient changes as the concentrations of reactants and products change until equilibrium is reached.



## 2. Law of Chemical Equilibrium (or law of mass action)

(a) For a reversible reaction represented by  $a A + b B \xrightleftharpoons[k_b]{k_f} c C + d D$ ,

At time = t, Rate of forward reaction,  $R_f = k_f [A]_t^a [B]_t^b$

Rate of backward reaction,  $R_b = k_b [C]_t^c [D]_t^d$

At equilibrium,  $R_f = R_b$

$\Rightarrow k_f [A]^a [B]^b = k_b [C]^c [D]^d$  (subscript 'eqm' is left out for convenience)

Hence, reaction quotient,  $Q_c = \frac{[C]^c [D]^d}{[A]^a [B]^b} = \frac{k_f}{k_b} = K_c$ , Equilibrium constant

(b) Relationship between,  $Q_c$ ,  $K_c$  and equilibrium

$Q_c < K_c$	$Q_c = K_c$	$Q_c > K_c$
System not at eqm. Forward rxn is favoured and $Q_c$ increases until $Q_c = K_c$ .	System at equilibrium	System not at eqm. Backward reaction is favoured and $Q_c$ decreases until $Q_c = K_c$ .

(c) Equilibrium constant is a constant at a fixed temperature. Hence, the value of equilibrium constant will not change unless temperature changes. The effect of temperature on the value of equilibrium constant is shown in the table below:

$\Delta H$ of forward reaction	$>0$ (endothermic)	$= 0$	$<0$ (exothermic)
Effect of $K_c$ when temperature $\uparrow$	$K_c \uparrow$	No change	$K_c \downarrow$
Effect of $K_c$ when temperature $\downarrow$	$K_c \downarrow$	No change	$K_c \uparrow$

### Types of Equilibrium Constants

Equilibrium Constant	Reactions	Example	Expression & Units
$K_c$ , involving concentrations in $\text{mol dm}^{-3}$	Reactions involving gases or aqueous solutions	$2\text{VO}^{2+}(\text{aq}) + \text{Cu}(\text{s}) + 4\text{H}^+(\text{aq}) \rightleftharpoons 2\text{V}^{3+}(\text{aq}) + \text{Cu}^{2+}(\text{aq}) + 2\text{H}_2\text{O}(\text{l})$	$K_c = \frac{[\text{V}^{3+}(\text{aq})]^2 [\text{Cu}^{2+}(\text{aq})]}{[\text{VO}^{2+}(\text{aq})]^2 [\text{H}^+(\text{aq})]^4}$
	Reactions involving liquids	$\text{CH}_3\text{COOH}(\text{l}) + \text{CH}_3\text{OH}(\text{l}) \rightleftharpoons \text{CH}_3\text{COOCH}_3(\text{l}) + \text{H}_2\text{O}(\text{l})$	$K_c = \frac{[\text{CH}_3\text{COOCH}_3(\text{l})][\text{H}_2\text{O}(\text{l})]}{[\text{CH}_3\text{COOH}(\text{l})][\text{CH}_3\text{OH}(\text{l})]}$
$K_p$ , involving partial pressure	Reactions involving gases	$3\text{H}_2(\text{g}) + \text{N}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$	$K_p = \frac{(P_{\text{NH}_3})^2}{(P_{\text{H}_2})^3 P_{\text{N}_2}}$

### Some points to take note when writing expressions for Equilibrium Constants

1. In general, for a reversible reaction represented by  $a A + b B \rightleftharpoons c C + d D$ ,

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b} \text{ or } K_p = \frac{P_C^c P_D^d}{P_A^a P_B^b} \text{ (for gaseous reactions only)}$$

- Leave out substances whose concentration is a constant (eg solids, immiscible liquids or water in reactions occurring in aqueous solutions) for  $K_c$  or whose partial pressure is very low and constant (eg solids or liquids) for  $K_p$ .

### Relationship between K and equation written

Expression for K is dependent on the equation written:

Equation	Expression for K	Relation to $K_1$
$3H_2(g) + N_2(g) \rightleftharpoons 2NH_3(g)$	$K_1 = \frac{[NH_3(g)]^2}{[H_2(g)]^3 [N_2(g)]}$	-----
$\frac{3}{2}H_2(g) + \frac{1}{2}N_2(g) \rightleftharpoons NH_3(g)$	$K_2 = \frac{[NH_3(g)]}{[H_2(g)]^{\frac{3}{2}} [N_2(g)]^{\frac{1}{2}}}$	$K_2 = \sqrt{K_1}$
$2NH_3(g) \rightleftharpoons 3H_2(g) + N_2(g)$	$K_3 = \frac{[H_2(g)]^3 [N_2(g)]}{[NH_3(g)]^2}$	$K_3 = (1 / K_1)$
$3H_2(g) + N_2(g) \rightleftharpoons 2NH_3(l)$	$K_4 = \frac{1}{[H_2(g)]^3 [N_2(g)]}$	-----

### Le Chatelier's Principle and Using it to Predict Shift in Position of Equilibrium

- Le Chatelier's Principle states that if a system in equilibrium is subjected to a change which disturbs the equilibrium, the system responds in such a way to counteract the effect of the change.
- Predict and explain the effect of the following changes on the yield of  $NH_3$  for the reaction:



No	Change	Effect of yield of $NH_3$	Reason
1	Adding more $N_2$	↑	Forward rxn favoured to decrease $[N_2]$ .
2	Using more iron catalyst	no change	The Fe catalyst affects the forward and backward reaction to the same extent.
3	Increasing temperature	↓	Backward reaction favoured to remove heat.
4	Higher pressure	↑	Forward rxn favoured as it gives less gaseous molecules so that pressure is decreased.
5	Using a reaction vessel of a larger volume	↓	↑ in volume decreases total concentration. Backward rxn favoured as it gives more gaseous molecules to ↑ total concentration.

### Problems on Equilibrium Involving Gases

- Useful formulae:
  - Partial pressure of a gas = mole fraction of gas x total pressure  

$$P_A = X_A P_{\text{total}} \text{ where } X_A = \frac{\text{no of mol of A}}{\text{total no of mol}} = \frac{n_A}{n_{\text{total}}}$$
  - Degree of dissociation of A,  $\alpha = \frac{\text{amt of A dissociated}}{\text{initial amt of A}}$

- (c) For a gaseous mixture made up of gases A and B with mole fraction of  $X_A$  and  $X_B$  respectively, (Note:  $X_B = 1 - X_A$ )  
Apparent  $M_r = X_A \times M_r \text{ of A} + X_B \times M_r \text{ of B}$   
When mass and total number of mol of a gaseous mixture is given, apparent  $M_r = \frac{\text{mass}}{n_{\text{total}}}$  ( $n_{\text{total}}$  is often found using  $pV = nRT$  equation)

## G. Acid Base Equilibria

### (I) Definitions (Know definitions of these terms):

Terms	Definition
Bronsted acid	Proton donor in aq. solution
Bronsted base	Proton acceptor in aq. solution
Strong acid/base	Fully dissociated Note: concentration of acid $\neq$ strength
Weak acid/base	Partially dissociated
pH	$= -\lg[H^+]$
pOH	$= -\lg[OH^-]$
$K_w$	$= [H^+][OH^-] = 1.0 \times 10^{-14} \text{ mol}^2\text{dm}^{-6}$ at 298 K
$K_a$	Acid dissociation constant, indicator of strength of acid. Higher $K_a \rightarrow$ stronger acid.  $HA(aq) \rightleftharpoons H^+(aq) + A^-(aq)$ $K_a = \frac{[H^+][A^-]}{[HA]} \text{ mol dm}^{-3}$
$pK_a$	$= -\lg K_a$
$pK_b$	Base dissociation constant, indicator of strength of base. Higher $K_b \rightarrow$ stronger base
$pK_b$	$= -\lg K_b$
$K_w$	$= K_a K_b$ applies only for acid-conjugate base pair

## (II) Calculations of pH for Different Types of Solutions

Solutions with pH < 7	Solutions with pH > 7																
<p>1. <b>Strong acid:</b> Calculate pH of a solution of dilute H<sub>2</sub>SO<sub>4</sub> with concentration of 0.025 mol dm<sup>-3</sup>.</p> <p>[H<sup>+</sup> (aq)] = 2 x 0.025 = 0.0500 mol dm<sup>-3</sup></p> <p>pH = -log [H<sup>+</sup> (aq)] = 1.30</p>	<p>1. <b>Strong base:</b> Calculate pH of a solution of aqueous Ba(OH)<sub>2</sub> with concentration of 0.025 mol dm<sup>-3</sup>.</p> <p>[OH<sup>-</sup> (aq)] = 2 x 0.025 = 0.0500 mol dm<sup>-3</sup></p> <p>pOH = -log [OH<sup>-</sup> (aq)] = 1.30; pH = 14 - pOH = 12.7</p>																
<p>2. <b>Weak acid, HA:</b> Calculate pH of a solution of aq CH<sub>3</sub>COOH with concentration of 0.050 mol dm<sup>-3</sup> given K<sub>a</sub> of CH<sub>3</sub>COOH is 1.8 x 10<sup>-5</sup> mol dm<sup>-3</sup></p> <p>Let [H<sup>+</sup> (aq)] at eqm be a mol dm<sup>-3</sup>. Ignoring H<sup>+</sup> from self-ionisation of water, [CH<sub>3</sub>COO<sup>-</sup> (aq)] = a mol dm<sup>-3</sup></p> <p>CH<sub>3</sub>COOH(aq) ⇌ CH<sub>3</sub>COO<sup>-</sup>(aq) + H<sup>+</sup>(aq)</p> <table border="1"> <tr> <td>Initial conc<sup>n</sup>/ mol dm<sup>-3</sup></td> <td>0.050</td> <td>–</td> <td>–</td> </tr> <tr> <td>Eqm conc<sup>n</sup>/ mol dm<sup>-3</sup></td> <td>0.050 - a</td> <td>a</td> <td>a</td> </tr> </table> <p>K<sub>a</sub> = <math>\frac{[\text{CH}_3\text{COO}^-][\text{H}^+]}{[\text{CH}_3\text{COOH}]}</math> = <math>\frac{a^2}{0.050 - a} \approx \frac{a^2}{0.050}</math> (since 0.050 &gt;&gt; a)</p> <p>[H<sup>+</sup> (aq)] = a ≈ √[K<sub>a</sub>[HA]<sub>0</sub>] = 9.487 x 10<sup>-4</sup> mol dm<sup>-3</sup></p> <p>pH = -log (9.487 x 10<sup>-4</sup>) = 3.02</p>	Initial conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050	–	–	Eqm conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050 - a	a	a	<p>2. <b>Weak base, B:</b> Calculate pH of a solution of aq NH<sub>3</sub> with concentration of 0.050 mol dm<sup>-3</sup> given K<sub>b</sub> of NH<sub>3</sub> is 1.74 x 10<sup>-5</sup> mol dm<sup>-3</sup></p> <p>Let [OH<sup>-</sup> (aq)] at eqm be a mol dm<sup>-3</sup>. Ignoring OH<sup>-</sup> from self-ionisation of water, [NH<sub>4</sub><sup>+</sup> (aq)] = a mol dm<sup>-3</sup></p> <p>H<sub>2</sub>O(l) + NH<sub>3</sub>(aq) ⇌ NH<sub>4</sub><sup>+</sup>(aq) + OH<sup>-</sup>(aq)</p> <table border="1"> <tr> <td>Initial conc<sup>n</sup>/ mol dm<sup>-3</sup></td> <td>0.050</td> <td>–</td> <td>–</td> </tr> <tr> <td>Eqm conc<sup>n</sup>/ mol dm<sup>-3</sup></td> <td>0.050 - a</td> <td>a</td> <td>a</td> </tr> </table> <p>K<sub>b</sub> = <math>\frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]}</math> = <math>\frac{a^2}{0.050 - a} \approx \frac{a^2}{0.050}</math> (since 0.050 &gt;&gt; a)</p> <p>[OH<sup>-</sup> (aq)] = a ≈ √[K<sub>b</sub>[B]<sub>0</sub>] = 9.327 x 10<sup>-4</sup> mol dm<sup>-3</sup></p> <p>pOH = -log (9.327 x 10<sup>-4</sup>) = 3.03;</p> <p>pH = 14 - pOH = 10.97</p>	Initial conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050	–	–	Eqm conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050 - a	a	a
Initial conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050	–	–														
Eqm conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050 - a	a	a														
Initial conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050	–	–														
Eqm conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050 - a	a	a														
SALT HYDROLYSIS CALCULATIONS																	
<p>3. <b>Salt of weak base, BH<sup>+</sup> (strong conjugate acid):</b> Calculate pH of a solution of aq NH<sub>4</sub>Cl with concentration of 0.050 mol dm<sup>-3</sup> given K<sub>b</sub> of NH<sub>3</sub> is 1.74 x 10<sup>-5</sup> mol dm<sup>-3</sup></p> <p>Let [H<sup>+</sup> (aq)] at eqm be a mol dm<sup>-3</sup>. Ignoring H<sup>+</sup> from self-ionisation of water, [NH<sub>3</sub> (aq)] = a mol dm<sup>-3</sup></p> <p>NH<sub>4</sub><sup>+</sup> (aq) ⇌ NH<sub>3</sub> (aq) + H<sup>+</sup>(aq)</p> <table border="1"> <tr> <td>Initial conc<sup>n</sup>/ mol dm<sup>-3</sup></td> <td>0.050</td> <td>–</td> <td>–</td> </tr> <tr> <td>Eqm conc<sup>n</sup>/ mol dm<sup>-3</sup></td> <td>0.050 - a</td> <td>a</td> <td>a</td> </tr> </table> <p>K<sub>a</sub> = <math>\frac{K_w}{K_b}</math> = <math>\frac{[\text{NH}_3][\text{H}^+]}{[\text{NH}_4^+]}</math> = <math>\frac{a^2}{0.050 - a} \approx \frac{a^2}{0.050}</math> (since 0.050 &gt;&gt; a)</p> <p>[H<sup>+</sup> (aq)] = a ≈ √<math>\frac{K_w}{K_b}</math>[BH<sup>+</sup>]<sub>0</sub>] = 5.361 x 10<sup>-6</sup> mol dm<sup>-3</sup></p> <p>pH = -log (5.361 x 10<sup>-6</sup>) = 5.27</p>	Initial conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050	–	–	Eqm conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050 - a	a	a	<p>3. <b>Salt of weak acid, A<sup>-</sup> (strong conjugate base):</b> Calculate pH of a solution of aq CH<sub>3</sub>COONa with concentration of 0.050 mol dm<sup>-3</sup> given K<sub>a</sub> of CH<sub>3</sub>COOH is 1.8 x 10<sup>-5</sup> mol dm<sup>-3</sup></p> <p>Let [OH<sup>-</sup> (aq)] at eqm be a mol dm<sup>-3</sup>. Ignoring OH<sup>-</sup> from self-ionisation of water, [CH<sub>3</sub>COOH (aq)] = a mol dm<sup>-3</sup></p> <p>H<sub>2</sub>O + CH<sub>3</sub>COO<sup>-</sup> ⇌ CH<sub>3</sub>COOH + OH<sup>-</sup></p> <table border="1"> <tr> <td>Initial conc<sup>n</sup>/ mol dm<sup>-3</sup></td> <td>0.050</td> <td>–</td> <td>–</td> </tr> <tr> <td>Eqm conc<sup>n</sup>/ mol dm<sup>-3</sup></td> <td>0.050 - a</td> <td>a</td> <td>a</td> </tr> </table> <p>K<sub>b</sub> = <math>\frac{K_w}{K_a}</math> = <math>\frac{[\text{CH}_3\text{COOH}][\text{OH}^-]}{[\text{CH}_3\text{COO}^-]}</math> = <math>\frac{a^2}{0.050 - a} \approx \frac{a^2}{0.050}</math> (since 0.050 &gt;&gt; a)</p> <p>[OH<sup>-</sup> (aq)] = a ≈ √<math>\frac{K_w}{K_a}</math>[A<sup>-</sup>]<sub>0</sub>] = 5.270 x 10<sup>-6</sup> mol dm<sup>-3</sup></p> <p>pOH = -log (5.270 x 10<sup>-6</sup>) = 5.28;</p> <p>pH = 14 - pOH = 8.72</p>	Initial conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050	–	–	Eqm conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050 - a	a	a
Initial conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050	–	–														
Eqm conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050 - a	a	a														
Initial conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050	–	–														
Eqm conc <sup>n</sup> / mol dm <sup>-3</sup>	0.050 - a	a	a														

### BUFFERS CALCULATIONS

4. **Acid buffer made up of weak acid and its salt, HA and A<sup>-</sup>:** Calculate pH of a mixture made of 20 cm<sup>3</sup> of 0.050 mol dm<sup>-3</sup> aq CH<sub>3</sub>COOH and 40 cm<sup>3</sup> of 0.030 mol dm<sup>-3</sup> aq CH<sub>3</sub>COONa.

(K<sub>a</sub> of CH<sub>3</sub>COOH = 1.8 × 10<sup>-5</sup> mol dm<sup>-3</sup>)

After mixing,

$$[\text{CH}_3\text{COOH}] = \frac{20}{20+40} \times 0.050 = 0.01667 \text{ mol dm}^{-3}$$

$$[\text{CH}_3\text{COO}^-] = \frac{40}{20+40} \times 0.030 = 0.020 \text{ mol dm}^{-3}$$

$$\text{pH} = \text{pK}_a + \log \frac{[\text{A}^-]}{[\text{HA}]}$$

$$= -\log(1.8 \times 10^{-5}) + \log \frac{0.020}{0.01667} = 4.82$$

4. **Alkaline buffer made of weak base and its salt, B and BH<sup>+</sup>:** Calculate pH of a mixture made of 20 cm<sup>3</sup> of 0.050 mol dm<sup>-3</sup> aq NH<sub>3</sub> and 40 cm<sup>3</sup> of 0.030 mol dm<sup>-3</sup> aq NH<sub>4</sub><sup>+</sup>.  
(K<sub>b</sub> of NH<sub>3</sub> = 1.74 × 10<sup>-5</sup> mol dm<sup>-3</sup>)

After mixing,

$$[\text{NH}_3] = \frac{20}{20+40} \times 0.050 = 0.01667 \text{ mol dm}^{-3}$$

$$[\text{NH}_4^+] = \frac{40}{20+40} \times 0.030 = 0.020 \text{ mol dm}^{-3}$$

$$\text{pOH} = \text{pK}_b + \log \frac{[\text{BH}^+]}{[\text{B}]}$$

$$= -\log(1.74 \times 10^{-5}) + \log \frac{0.020}{0.01667} = 4.84$$

$$\text{pH} = 14 - \text{pOH} = 9.16$$

### (III) Explanation of how buffer works using equations:



Acid Buffer: made up of weak acid HA and its salt NaA	Alkaline Buffer: made up of weak base B and its salt (eg BH <sup>+</sup> Cl <sup>-</sup> )
<p><u>On addition of H<sup>+</sup>:</u>  <math>\text{A}^- (\text{aq}) + \text{H}^+ (\text{aq}) \longrightarrow \text{HA} (\text{aq})</math></p> <p style="text-align: center;">Single arrow to show removal of H<sup>+</sup></p> <p>A<sup>-</sup> present reacts with H<sup>+</sup> so that [H<sup>+</sup>] remains approximately constant and pH is maintained.</p>	<p><u>On addition of H<sup>+</sup>:</u>  <math>\text{B} (\text{aq}) + \text{H}^+ (\text{aq}) \longrightarrow \text{BH}^+ (\text{aq})</math></p> <p>B present reacts with H<sup>+</sup> so that [H<sup>+</sup>] remains approximately constant and pH is maintained.</p>
<p><u>On addition of OH<sup>-</sup>:</u>  <math>\text{HA} (\text{aq}) + \text{OH}^- (\text{aq}) \longrightarrow \text{A}^- (\text{aq}) + \text{H}_2\text{O} (\text{l})</math></p> <p>HA present reacts with OH<sup>-</sup> so that [OH<sup>-</sup>] remains approximately constant and pH is maintained.                      Large reservoir of HA and A<sup>-</sup> present is able to cope with small amount of H<sup>+</sup> and OH<sup>-</sup> added.</p>	<p><u>On addition of OH<sup>-</sup>:</u>  <math>\text{BH}^+ (\text{aq}) + \text{OH}^- (\text{aq}) \longrightarrow \text{B} (\text{aq}) + \text{H}_2\text{O} (\text{l})</math></p> <p>BH<sup>+</sup> present reacts with OH<sup>-</sup> so that [OH<sup>-</sup>] remains approximately constant and pH is maintained.                      Large reservoir of B and BH<sup>+</sup> present is able to cope with small amount of H<sup>+</sup> and OH<sup>-</sup> added.</p>

### (IV) Application: Role of H<sub>2</sub>CO<sub>3</sub>/HCO<sub>3</sub><sup>-</sup> in controlling pH of blood

Buffer system in human blood:

On addition of H<sup>+</sup>:

On addition of OH<sup>-</sup>:



### Regions in Titration Curve

Titration Curve of Weak Acid vs Strong Base (WA-SB)			Titration Curve of Weak Base vs Strong Acid (WB-SA)		
Region	①	②	Region	①	②
Contents	Weak acid, HA	Mixture of HA and A <sup>-</sup>	Contents	Weak base, B	Mixture of B & HB <sup>+</sup>
[H <sup>+</sup> ] or pH	$[H^+] \approx \sqrt{K_a[HA]_0}$	$pH = pK_a + \lg \frac{[A^-]}{[HA]}$ $= pK_a + \lg \frac{n_{OH^- added}}{n_{HA^0} - n_{OH^- added}}$  At $\frac{1}{2}V_E$ : $pH = pK_a$ (max buffer capacity)	[OH <sup>-</sup> ] or pOH	$[OH^-] \approx \sqrt{K_b[B]_0}$	$pOH = pK_b + \lg \frac{[BH^+]}{[B]}$ $= pK_b + \lg \frac{n_{H^+ added}}{n_{B^0} - n_{H^+ added}}$  At $\frac{1}{2}V_E$ : $pH = pK_a = 14 - pK_b$ (max buffer capacity)
Region	③ (equivalence pt)	④	Region	③ (equivalence pt)	④
Contents	Salt of weak acid, A <sup>-</sup>	Excess base	Contents	Salt of weak base, BH <sup>+</sup>	Excess acid
[OH <sup>-</sup> ] or pOH	$[OH^-] \approx \sqrt{\frac{K_w}{K_a}[A^-]_0}$	$[OH^-] = \frac{\text{amt of xs OH}}{\text{Vol}}$	[H <sup>+</sup> ] or pH	$[H^+] \approx \sqrt{\frac{K_w}{K_b}[BH^+]_0}$	$[H^+] = \frac{\text{amt of xs H}^+}{\text{Vol}}$

Note: you should still show clear and proper workings and not just apply these formulae blindly.

## H. Solubility Equilibria

### (I) $K_{sp}$ and solubility

General Example - $A_bB_a$ made up of $A^{a+}$ & $B^{b-}$	Specific Example - $Ca_3(PO_4)_2$
1. For a sparingly soluble compound $A_bB_a$ : $A_bB_a(s) + aq \rightleftharpoons bA^{a+}(aq) + aB^{b-}(aq)$ ; $K_{sp} = [A^{a+}]^b[B^{b-}]^a$	Eg calcium phosphate: $Ca_3(PO_4)_2$ $Ca_3(PO_4)_2(s) + aq \rightleftharpoons 3Ca^{2+}(aq) + 2PO_4^{3-}(aq)$ ; $K_{sp} = [Ca^{2+}]^3[PO_4^{3-}]^2$
2. If solubility of $A_bB_a$ in water is $s \text{ mol dm}^{-3}$ , $[A^{a+}] = bs \text{ mol dm}^{-3}$ , $[B^{b-}] = as \text{ mol dm}^{-3}$	If solubility of $Ca_3(PO_4)_2$ is $2.6 \times 10^{-6} \text{ mol dm}^{-3}$ , $[Ca^{2+}(aq)] = 3 \times 2.6 \times 10^{-6} = 7.8 \times 10^{-6} \text{ mol dm}^{-3}$ $[PO_4^{3-}(aq)] = 2 \times 2.6 \times 10^{-6} = 5.2 \times 10^{-6} \text{ mol dm}^{-3}$
3. If solubility of $A_bB_a$ in water is $s \text{ mol dm}^{-3}$ , $K_{sp} = [A^{a+}]^b[B^{b-}]^a = (bs)^b(as)^a = b^b a^a s^{(a+b)}$	If solubility of $Ca_3(PO_4)_2$ is $2.6 \times 10^{-6} \text{ mol dm}^{-3}$ , $K_{sp} = [Ca^{2+}]^3[PO_4^{3-}]^2$ $= (3 \times 2.6 \times 10^{-6})^3 (2 \times 2.6 \times 10^{-6})^2$ $= 1.3 \times 10^{-26} \text{ mol}^5 \text{ dm}^{-15}$
4. If solubility product of $A_bB_a$ in water is $K_{sp}$ , then its solubility $s = \sqrt[a+b]{\frac{K_{sp}}{b^b a^a}}$	Given $K_{sp}$ of $Ca_3(PO_4)_2 = 1.3 \times 10^{-26} \text{ mol}^5 \text{ dm}^{-15}$ , calculate its solubility. Let its solubility be $s \text{ mol dm}^{-3}$ $K_{sp} = [Ca^{2+}]^3[PO_4^{3-}]^2 = (3s)^3(2s)^2 = 108 s^5$ $s = \sqrt[5]{\frac{K_{sp}}{3^3 2^2}} = \sqrt[5]{\frac{1.3 \times 10^{-26}}{3^3 2^2}} = 2.61 \times 10^{-6} \text{ mol dm}^{-3}$

### (II) Ionic Product, $K_{sp}$ and Precipitation

1. Precipitation occurs when ionic product is greater than  $K_{sp}$ .

Ionic product = $[M^+(aq)][X^-(aq)]$	Remark	Precipitation?
$IP < K_{sp}$	The solution is unsaturated.	No!
$IP = K_{sp}$	The solution is saturated.	No
$IP > K_{sp}$	The solution is beyond saturation upon mixing.	Yes!

Note: Ionic product and  $K_{sp}$  have the same expression. The only difference is for IP, the concentration terms are the concentration at a particular instant, but for  $K_{sp}$ , equilibrium concentrations are used.

2. To predict whether precipitation occurs when two solutions, one containing cations and the other containing anions of a sparingly soluble salt, calculate
  - (i) the new concentration of the cation and anion immediately after mixing
  - (ii) ionic product
If ionic product  $> K_{sp}$ , then precipitation occurs.

### (III) Change in solubility of a sparingly soluble salt

1. The solubility of a sparingly soluble salt is decreased in the presence of its cation or anion. This is known as the common ion effect. For example, solubility of silver iodide is decreased in the presence of silver ions or iodide ions.
2. The solubility of a sparingly soluble salt may be increased by adding a reagent which reacts with its cation or anion. The phenomenon can either be explained using Le Chatelier's Principle or ionic product. Some examples:



Sparingly soluble salt	Reagent added to increase its solubility	Remarks
Silver chloride $\text{AgCl (s)} + \text{aq} \rightleftharpoons \text{Ag}^+(\text{aq}) + \text{Cl}^-(\text{aq})$	Aq $\text{NH}_3$ $\text{Ag}^+(\text{aq}) + 2\text{NH}_3(\text{aq}) \rightleftharpoons \text{Ag}(\text{NH}_3)_2^+(\text{aq})$	$K_{\text{sp}}$ of AgCl is not so low so that it is soluble in xs aq $\text{NH}_3$ . However, AgBr and AgI are insoluble in xs aq $\text{NH}_3$ as their $K_{\text{sp}}$ are too low.
Barium chromate: $\text{BaCrO}_4(\text{s}) + \text{aq} \rightleftharpoons \text{Ba}^{2+}(\text{aq}) + \text{CrO}_4^{2-}(\text{aq})$	Dilute nitric acid $2\text{CrO}_4^{2-}(\text{aq}) + 2\text{H}^+(\text{aq}) \rightleftharpoons \text{Cr}_2\text{O}_7^{2-}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	$\text{Ag}_2\text{CrO}_4$ also dissolves in dil $\text{HNO}_3$ due to the reaction between $\text{CrO}_4^{2-}$ and $\text{H}^+$ .
Copper(II) hydroxide $\text{Cu}(\text{OH})_2(\text{s}) + \text{aq} \rightleftharpoons \text{Cu}^{2+}(\text{aq}) + 2\text{OH}^-(\text{aq})$	Aq $\text{NH}_3$ $\text{Cu}^{2+}(\text{aq}) + 4\text{NH}_3(\text{aq}) \rightleftharpoons \text{Cu}(\text{NH}_3)_4^{2+}(\text{aq})$  $[\text{Cu}^{2+}]$ decreases due to formation of $\text{Cu}(\text{NH}_3)_4^{2+}$ complex $\rightarrow$ equilibrium position for $\text{Cu}(\text{OH})_2(\text{s}) + \text{aq} \rightleftharpoons \text{Cu}^{2+}(\text{aq}) + 2\text{OH}^-(\text{aq})$ shifts right $\rightarrow$ ppt $\text{Cu}(\text{OH})_2(\text{s})$ dissolve	Zinc hydroxide also dissolves in excess aq $\text{NH}_3$ due to the formation of $\text{Zn}(\text{NH}_3)_4^{2+}(\text{aq})$

## I. Electrochemistry

Assessment Objectives	Remarks
(a) describe and explain redox processes in terms of electron transfer and/or of changes in oxidation number (oxidation state).	Oxidation = loss of electrons/ $\uparrow$ in oxidation state Reduction = gain of electrons/ $\downarrow$ in oxidation state
(b) define the terms: (i) standard electrode (redox) potential, $E^\ominus$ (ii) standard cell potential, $E^\ominus_{\text{cell}}$	$E^\ominus$ of a standard half-cell is defined as the potential of that half-cell relative to a standard hydrogen half-cell, under standard conditions. $E^\ominus_{\text{cell}}$ is the potential difference between 2 half-cells under standard conditions
(c) describe the standard hydrogen electrode.	Draw the experimental set-up to help you in your description.
(d) describe methods used to measure the standard electrode potentials of (i) metals or non-metals in contact with their ions in aqueous solution (ii) ions of the same element in different oxidation states	
(e) calculate a standard cell potential by combining two standard electrode potentials.	$E^\ominus_{\text{cell}} = E^\ominus_{\text{Red}} - E^\ominus_{\text{Ox}}$ (where $E^\ominus_{\text{Red}} = E^\ominus$ of substance reduced and $E^\ominus_{\text{Ox}} = E^\ominus$ of substance oxidised)
(f) use standard cell potentials to (i) explain/predict the direction of electron flow from a simple cell; (ii) predict the feasibility of a reaction	(i) $\frac{1}{2}$ -cell with bigger $E^\ominus$ is the [R] $\frac{1}{2}$ -cell and hence cathode. The other $\frac{1}{2}$ -cell with smaller $E^\ominus$ is the [O] $\frac{1}{2}$ -cell and hence anode. Electron flows from anode to cathode. (ii) In predicting feasibility of a reaction, first identify relevant $\frac{1}{2}$ -eqn and compute $E^\ominus_{\text{cell}}$ . Reaction is energetically feasible if $E^\ominus_{\text{cell}} > 0$
(g) Understand the limitations in the use of standard cell potentials to predict the feasibility of a reaction.	Limitations = rxn may be kinetically slow, conditions not std and side rxn may occur.
(h) construct redox equations using the relevant half-equations (including the use of $\text{Fe}^{3+} / \text{Fe}^{2+}$ , $\text{MnO}_4^- / \text{Mn}^{2+}$ and $\text{Cr}_2\text{O}_7^{2-} / \text{Cr}^{3+}$ as examples of redox systems).	$\text{MnO}_4^-$ reduced to $\text{Mn}^{2+}$ under acidified medium and to $\text{MnO}_2$ under weakly acidic/neutral/ weakly alkaline conditions. Under strongly alkaline conditions, $\text{MnO}_4^-$ is reduced to $\text{MnO}_4^{2-}$ .
(i) predict qualitatively how the value of an electrode potential varies with the concentration of the aqueous ion.	Changes that $\uparrow E_{\text{Red}}$ or $\downarrow E_{\text{Ox}}$ increase $E_{\text{cell}}$ . Changes that $\downarrow E_{\text{Red}}$ or $\uparrow E_{\text{Ox}}$ decrease $E_{\text{cell}}$ .

(j) state the possible advantages of developing other types of cell, e.g. the $\text{H}_2/\text{O}_2$ fuel cell and improved batteries (as in electric vehicles) in terms of smaller size, lower mass and higher voltage.	Learn how to draw $\text{H}_2/\text{O}_2$ fuel cell and write equations for the reactions at anode/cathode.
(k) state the relationship between the Faraday constant, the Avogadro constant and the charge on the electron, $F = Le$ .	–
(l) predict the identity of the substance liberated during electrolysis from the state of electrolyte (molten or aqueous), position in the redox series (electrode potential) and concentration.	Zinc and metals more reactive than it cannot be obtained from electrolysis of their aqueous solutions. Water is reduced instead: $2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2 + 2\text{OH}^-$ $\text{Cl}^-$ is oxidised at the anode if concentrated solution is used; $\text{I}^-$ and $\text{Br}^-$ are oxidised at anode from aq solutions. Sulfate and nitrate ions are not discharged. Instead water is oxidised: $2\text{H}_2\text{O} \rightarrow \text{O}_2 + 4\text{H}^+ + 4\text{e}^-$
(m) calculate: (i) the quantity of charge passed during electrolysis; (ii) the mass and/or volume of substance liberated during electrolysis, including those in the electrolysis of $\text{H}_2\text{SO}_4$ (aq), $\text{Na}_2\text{SO}_4$ (aq).	$Q = I \times t$ or $Q = nF$ where $Q$ = quantity of charge in coulombs $I$ = current in amperes $t$ = time in seconds $F$ = Faraday's Constant
(n) explain, in terms of the electrode reactions, the industrial processes of: (i) the anodising of aluminium; (ii) the electrolytic purification of copper.	Learn description and how to draw experimental set up as well.

### Equations showing discharge of $\text{H}_2$ and $\text{O}_2$ from aqueous solutions

Discharge of $\text{H}_2$ at cathode		Discharge of $\text{O}_2$ at anode	
Type of aq solution electrolysed	Equation	Type of aq solution electrolysed	Equation
Acidic eg dil $\text{H}_2\text{SO}_4$	$2\text{H}^+ (\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2$	Acidic eg dil $\text{H}_2\text{SO}_4$	$2\text{H}_2\text{O} (\text{l}) \rightarrow \text{O}_2(\text{g}) + 4\text{H}^+ (\text{aq}) + 4\text{e}^-$
Neutral eg aq $\text{Na}_2\text{SO}_4$	$2\text{H}_2\text{O} (\text{l}) + 2\text{e}^- \rightarrow \text{H}_2 (\text{g}) + 2\text{OH}^- (\text{aq})$	Neutral eg aq $\text{Na}_2\text{SO}_4$	$2\text{H}_2\text{O} (\text{l}) \rightarrow \text{O}_2(\text{g}) + 4\text{H}^+ (\text{aq}) + 4\text{e}^-$
Alkaline eg aq $\text{KOH}$	$2\text{H}_2\text{O} (\text{l}) + 2\text{e}^- \rightarrow \text{H}_2 (\text{g}) + 2\text{OH}^- (\text{aq})$	Alkaline eg aq $\text{KOH}$	$4\text{OH}^- (\text{aq}) \rightarrow \text{O}_2(\text{g}) + 2\text{H}_2\text{O} (\text{l}) + 4\text{e}^-$



# RI YEAR 5 – 6 ORGANIC CHEMISTRY SUMMARY

## A Formulae

Types of Formulae	Compound	2-methylpent-1-ene	4-cyanocyclohexanol	3-ethyl-4-nitrobenzoic acid
<b>Empirical Formula</b> (simplest formula)		CH <sub>2</sub>	C <sub>7</sub> H <sub>11</sub> ON	C <sub>9</sub> H <sub>9</sub> O <sub>4</sub> N
<b>Molecular Formula</b> (shows no. of atoms of each element)		C <sub>6</sub> H <sub>12</sub>	C <sub>7</sub> H <sub>11</sub> ON	C <sub>9</sub> H <sub>9</sub> O <sub>4</sub> N
<b>Structural Formula</b> (shows arrangement of atoms without ambiguity)		$\begin{array}{c} \text{CH}_3 \\   \\ \text{H}_2\text{C}=\text{C}-\text{CH}_2\text{CH}_2\text{CH}_3 \\   \\ \text{H} \end{array}$		
<b>Displayed Formula/ Full Structural Formula</b> (shows all bonds and atoms in open chains but for rings, symbols such as are acceptable)	 			
<b>Skeletal Formula</b> (Straight lines represent the C-C bonds. C & H atoms are not explicitly shown.)				

## B Hybridisation, Resonance and Delocalisation

Hybridisation		Delocalisation and Resonance (for species/ compounds ≠ metal)	
Hybridisation = mixing of orbitals resulting in hybrid orbitals formed. Each hybrid orbital has a big lobe and a small lobe.  Hybridisation allows geometries such as linear, trigonal planar, tetrahedral, trigonal bipyramidal and octahedral to be possible thus minimising repulsion between electron pairs. Three types of hybrid orbitals are encountered in H <sub>2</sub> chem:		Delocalisation = movement of electrons beyond the attraction of 3 or more p/d orbitals (⇒ π bond (s) present overlaps of 3 or more p/d orbitals (⇒ π bond (s) present in species with delocalisation of e <sup>-</sup> ) eg benzene Resonance = same attachment of atoms but different arrangement of electrons, eg benzene	
Type	Remarks	Shape & Diagram	
sp <sup>3</sup>	4 hybrid orbitals. No unhybridised p orbital ⇒ no π bond. For an atom with 4 regions of electron density.	Tetrahedral 	1. Species with delocalised electrons exhibit resonance and species that exhibit resonance have delocalised electrons.
sp <sup>2</sup>	3 hybrid orbitals. 1 unhybridised p orbital ⇒ may form 1 π bond. For an atom with 3 regions of electron density.	Trigonal planar 	2. How to recognise species which have delocalisation of electrons/resonance: (a) Species contain conjugated double bonds (ie at least two double bonds alternating with single bonds) <b>OR</b> (b) 'Bond Structure' diagrams of species can be drawn in more than 1 way
sp	2 hybrid orbitals. 2 unhybridised p orbitals ⇒ may form 2 π bonds. For an atom with 2 regions of electron density.	Linear 	3. Which of the following species contain delocalised electrons/ show resonance? <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;"> <p>(i)</p> </div> <div style="text-align: center;"> <p>(ii)</p> </div> <div style="text-align: center;"> <p>(iii)</p> </div> <div style="text-align: center;"> <p>(iv)</p> </div> <div style="text-align: center;"> <p>(v)</p> </div> <div style="text-align: center;"> <p>(vi)</p> </div> <div style="text-align: center;"> <p>(vii)</p> </div> </div>

## C Isomerism

### Isomerism


**Structural isomerism**  
(exhibited by substances with the same molecular formula but different structural formulae)

**Stereoisomerism**  
(exhibited by substances with same structural formula but different spatial arrangement of atoms.)

#### Functional Group Isomerism

eg  
 $\text{CH}_3\text{CH}_2\text{OH}$  &  $\text{CH}_3\text{OCH}_3$

#### Positional Isomerism



#### Straight chain vs branching

eg  
 $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$  &  $\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$

#### Geometric (Cis-trans) Isomerism


(caused by restriction of rotation due to double bond or ring.)

#### Optical Isomerism


(present in molecules with no plane of symmetry)

#### Chiral Carbon


Chiral carbon is  $\text{sp}^3$  hybridised and has four different groups attached to it. Mark the chiral carbons in the molecule shown below with an "\*" (there are eight):




Found in alkenes when the two groups attached to each of the two carbons in the double bond are different.



Found in rings when the two groups attached to two ring carbon atoms are different.




Compounds with 2 or more chiral carbons may not exhibit optical isomerism if a plane of symmetry is present in the molecule:



Such compounds are called meso compounds.

Compounds with only one chiral carbon exhibit optical isomerism:






Optical isomerism = existence of 2 cpds with the same structural formula but differ in being non-superimposable mirror images of each other.  
Geometric Isomerism = existence of 2 cpds with the same structural formula but have different spatial arrangement eg due to restricted rotation about a double bond or ring structure.

A compound that exhibits optical isomerism has two enantiomers (mirror images which are non-superimposable). If only one enantiomer is present, the compound is optically active and rotates plane-polarised light. If both enantiomers are present but in unequal proportions, the mixture still rotates plane-polarised light but the angle of rotation is smaller than that of the pure enantiomer. If both the enantiomers are present in equal proportions, the mixture is called a racemic mixture and shows no optical activity (ie does not rotate plane-polarised light).

## D Mechanism

**Free Radical Substitution in Alkanes:** eg Bromination of ethylbenzene. Alkane reacted with Br<sub>2</sub> in uv light.







Electrophilic Addition in Alkenes	Electrophilic Substitution in Arenes	Nucleophilic Substitution in Haloalkanes	Nucleophilic Addition in Carbonyl Compounds
<p>eg Bromination of propene</p> <p>Alkene reacted with bromine in CCl<sub>4</sub> or bromine water</p> <p><b>Addition of electrophile - formation of carbocation (Markovnikov's Rule)</b></p> $\text{CH}_3-\text{CH}=\text{CH}_2 \xrightarrow{\text{Br}_2} \text{CH}_3-\text{CH}(\text{Br})-\text{CH}_2\text{Br}$ <p><b>Addition of nucleophile:</b></p> $\text{CH}_3-\text{CH}(\text{Br})-\text{CH}_2\text{Br} \xrightarrow{\text{OH}^-} \text{CH}_3-\text{CH}(\text{OH})-\text{CH}_2\text{Br}$ <p>If bromine water is used, H<sub>2</sub>O can act as a nucleophile and bromohydrin is formed as the major product:</p> $\text{CH}_3-\text{CH}(\text{Br})-\text{CH}_2\text{Br} \xrightarrow{\text{H}_2\text{O}} \text{CH}_3-\text{CH}(\text{OH})-\text{CH}_2\text{Br}$	<p>eg Bromination of ethylbenzene</p> <p>Arene reacted with bromine in the presence of FeBr<sub>3</sub>.</p> <p><b>Generation of electrophile:</b></p> $\text{Br}_2 + \text{FeBr}_3 \rightarrow \text{Br}^+ + \text{FeBr}_4^-$ <p><b>Addition into benzene ring</b> (position dependent on group which is already present): <math>\text{CH}_2\text{CH}_3</math></p> <p></p> <p><b>Loss of H<sup>+</sup>:</b></p> <p></p> <p>+ HBr + FeBr<sub>3</sub></p>	<p>eg KCN and 2-chlorobutane</p> <p>Haloalkane warmed with KCN dissolved in alcohol.</p> <p>Since chloroethane is a secondary RX, the reaction can proceed via S<sub>N</sub>2 or S<sub>N</sub>1</p> <p><b>S<sub>N</sub>2 mechanism:</b></p> $\text{CH}_3\text{CH}_2\text{Cl} + \text{CN}^- \rightarrow [\text{CH}_3\text{CH}_2\text{Cl}\cdots\text{CN}]^\ddagger \rightarrow \text{CH}_3\text{CH}_2\text{CN} + \text{Cl}^-$ <p>Product from S<sub>N</sub>2 is optically active if reactant RX is optically active.</p> <p><b>S<sub>N</sub>1 mechanism:</b></p> $\text{CH}_3\text{CH}_2\text{Cl} \xrightarrow{\text{slow}} \text{CH}_3\text{CH}_2^+ + \text{Cl}^- \xrightarrow{\text{fast}} \text{CH}_3\text{CH}_2\text{CN}$ <p>Racemic mixture</p>	<p>eg HCN and butanone</p> <p>Aldehyde or ketone reacted with HCN with a small amount of base (or KCN).</p> <p><b>Generation of nucleophile:</b></p> $\text{HCN} + \text{OH}^- \rightarrow \text{CN}^- + \text{H}_2\text{O} \quad (\text{for NaOH})$ $\text{or } \text{KCN} \rightarrow \text{K}^+ + \text{CN}^- \quad (\text{for KCN})$ <p><b>Addition of CN<sup>-</sup>:</b></p> $\text{CH}_3\text{COCH}_2\text{CH}_3 + \text{CN}^- \xrightarrow{\text{slow}} [\text{CH}_3\text{C}(\text{O}^-)(\text{CN})\text{CH}_2\text{CH}_3]^\ddagger \xrightarrow{\text{fast}} \text{CH}_3\text{C}(\text{OH})(\text{CN})\text{CH}_2\text{CH}_3 + \text{CN}^-$ <p><b>Extraction of H<sup>+</sup> from HCN:</b></p> <p>The carbonyl C is sp<sup>2</sup> hybridised. CN<sup>-</sup> can add to either face with equal likelihood so that the resultant product forms a racemic mixture. Hence even though the product has one chiral carbon, it shows no optical activity.</p>

## E Acidity

1. A weak acid dissociates incompletely in water:  $\text{HA (aq)} \rightleftharpoons \text{H}^+ \text{(aq)} + \text{A}^- \text{(aq)}$ . The more stable the anion, the greater is the dissociation and hence the stronger is the acid.

2. In general,




Compound	Alcohol, ROH	Phenol, 	Carboxylic acid, RCOOH
Anion	Alkoxide, RO <sup>-</sup>	Phenoxide, 	Carboxylate ion, RCOO <sup>-</sup>
Stability of anion and relative acidity	The alkyl group is electron-donating so that the negative charge on O in RO <sup>-</sup> is more intensified compared to OH <sup>-</sup> . Hence RO <sup>-</sup> is destabilised by the electron-donating R group so that alcohols are weaker acids than water.	The p-orbital of O overlaps with the $\pi$ -electron cloud of the benzene ring so that the negative charge on O delocalises into the benzene ring. The dispersal of negative charge stabilises the phenoxide ion so that phenol is a stronger acid than water.	Carboxylate ion has 2 resonance structures: $\text{R}-\text{C}(=\text{O})-\text{O}^- \rightleftharpoons \text{R}-\text{C}(\text{O}^-)=\text{O}$ The 2 resonance structures are equivalent and negative charge is delocalised over two electronegative O atoms resulting in the carboxylate ions being greatly stabilised so that carboxylic acid is more acidic than phenol.
Reaction with Na	Effervescence of hydrogen: $\text{ROH (l)} + \text{Na (s)} \longrightarrow \text{RO}^-\text{Na}^+ \text{(s)} + \frac{1}{2}\text{H}_2 \text{(g)}$	Vigorous rxn with evolution of hydrogen: $\text{C}_6\text{H}_5\text{OH} + \text{Na} \longrightarrow \text{C}_6\text{H}_5\text{O}^-\text{Na}^+ \text{(s)} + \frac{1}{2}\text{H}_2 \text{(g)}$	Explosive rxn with evolution of hydrogen: $\text{RCOOH} + \text{Na} \longrightarrow \text{RCOO}^-\text{Na}^+ \text{(s)} + \frac{1}{2}\text{H}_2 \text{(g)}$
Rxn with aq NaOH	No reaction. Alcohol is not acidic enough to react with OH <sup>-</sup> .	Neutralisation reaction and phenol dissolves (to form 1 miscible layer) $\text{C}_6\text{H}_5\text{OH (s)} + \text{NaOH (aq)} \rightarrow \text{C}_6\text{H}_5\text{O}^-\text{Na}^+ \text{(aq)} + \text{H}_2\text{O (l)}$	Neutralisation reaction: $\text{RCOOH (aq)} + \text{NaOH (aq)} \rightarrow \text{RCOO}^-\text{Na}^+ \text{(aq)} + \text{H}_2\text{O (l)}$
Rxn with aq Na <sub>2</sub> CO <sub>3</sub>	No reaction. Alcohol is not acidic enough to react with CO <sub>3</sub> <sup>2-</sup> .	No reaction. Phenol is not acidic enough to react with CO <sub>3</sub> <sup>2-</sup> .	Evolution of CO <sub>2</sub> : $2\text{RCOOH} + \text{Na}_2\text{CO}_3 \rightarrow 2\text{RCOO}^-\text{Na}^+ + \text{H}_2\text{O} + \text{CO}_2 \text{(g)}$
Variation of acidity within class of compound	<p>Primary acid &gt; Secondary acid &gt; Tertiary alcohol</p> <p>eg</p> $\text{CH}_3\text{CH}_2\text{CH}_2-\text{C}(\text{H})_2-\text{OH} > \text{CH}_3\text{CH}_2-\text{C}(\text{H})(\text{CH}_3)-\text{OH} > \text{CH}_3-\text{C}(\text{H})(\text{CH}_3)_2-\text{OH}$ $\text{CH}_3\text{CH}_2\text{CH}_2-\text{C}(\text{H})_2-\text{O}^- > \text{CH}_3\text{CH}_2-\text{C}(\text{H})(\text{CH}_3)-\text{O}^- > \text{CH}_3-\text{C}(\text{H})(\text{CH}_3)_2-\text{O}^-$	<p>e<sup>-</sup>-donating groups, if present in phenol, reduce the delocalisation of negative charge of O into benzene ring thus decreasing the stability of phenoxide ion and its acidity. Conversely, e<sup>-</sup>-withdrawing groups enhance the delocalisation of negative charge of O into benzene ring thus increasing the stability of phenoxide ion and its acidity.</p> <p>eg</p> 	<p>An electron-donating group intensifies the -ve charge and decreases the acidity of a carboxylic acid while an electron-withdrawing group disperses the -ve charge and increases its acidity (same as phenol).</p> <p>Eg</p> $\text{CH}_3\text{COOH} < \text{CH}_2\text{ClCOOH} < \text{CH}_2\text{FCOOH}$ $\text{CH}_2\text{FCH}_2\text{COOH} < \text{CH}_3\text{CHFCOOH}$







## F Basicity

1. A base is a lone-pair donor. The more available the lone pair of electrons is for donation, the stronger is the base.

2. In general,










Compound	Amide, $\text{R}-\text{C}(=\text{O})-\text{NH}_2$	Phenylamine, 	Aliphatic amine, $\text{RNH}_2$
Availability of lone pair of electrons on N	<p>Amide has 2 resonance structures:</p> $\text{R}-\text{C}(=\text{O})-\text{NH}_2 \leftrightarrow \text{R}-\text{C}(\text{O}^-)=\text{NH}_2^+$ <p>Due to resonance, the lone pair of <math>e^-</math> on N is less available for protonation so that amide is neutral and less basic than phenylamine.</p> <p>Another explanation: p-orbital of N overlap with p-orbitals of carbonyl group resulting in delocalisation thus making lp on N less available.</p>	<p>The p-orbital of N overlaps with the <math>\pi</math>-electron cloud of the benzene ring so that the lone pair of electrons on N delocalise into the benzene ring. This decreases the availability of the lone pair of electrons on N for donation to a <math>\text{H}^+</math> (protonation). Hence phenylamine is a weaker base than ammonia.</p>	<p>Aliphatic amines contain alkyl groups. The alkyl group is electron donating and enhances the availability of the lone pair for protonation so that aliphatic amines are stronger bases than ammonia.</p> <p style="text-align: center;">  </p>
Variation of basicity within class of compound	—	<p><math>e^-</math>-donating groups, if present in phenylamine, reduce the delocalisation of lone pair of <math>e^-</math> on N into benzene ring thus increasing the availability of the lone pair of <math>e^-</math> for protonation thus increasing its basicity.</p> <p>Conversely, <math>e^-</math>-withdrawing groups enhance the delocalisation of lone pair of <math>e^-</math> on N into benzene ring thus decreasing availability of the lone pair of <math>e^-</math> for protonation thus decreasing its basicity.</p> <p>eg </p>	<p>As no of alkyl groups on N <math>\uparrow</math>, the availability of lone pair <math>e^-</math> for protonation is even more enhanced so that:</p> <p style="text-align: center;"> <math>\boxed{\text{Primary amine}} &lt; \boxed{\text{Secondary amine}} &lt; \boxed{\text{base}}</math> </p> <p>eg </p> <p>Tertiary amines are weaker bases than primary amines as the three bulky alkyl groups cause the lone pair of electrons on N to be less available due to steric hindrance.</p>




## G Physical Properties of Organic Compounds (Rough Guide, may vary due to branching of alkyl groups, class of alcohol, amine or amide etc)

Type of Organic Cpd	Hydrocarbon (alkane, alkene & arene)	Halogeno-alkanes	Alcohol	Phenol	Carbonyl Cpd (aldehyde & ketone)	Carboxylic acid	Ester	Amine, phenylamine	Amide
Dominant IMF	Mostly id-id interactions	pd-pd interactions	Hydrogen bonding	Hydrogen bonding	pd-pd interactions	Hydrogen bonding	pd-pd interactions	H bonding except 3° amine	Hydrogen bonding
State at rm temp	$\leq 4 \text{ C} = \text{gas}$ $> 5 \text{ C} = \text{liquid}$	Mostly liquids	$\leq 6 \text{ C} = \text{liquid}$ $> 7 \text{ C} = \text{solid}$	solid	Mostly liquids	$\leq 6 \text{ C} = \text{liquid}$ $> 7 \text{ C} = \text{solid}$	Mostly liquids	Mostly liquids	$\text{HCONH}_2 (\text{l})$ $> 2 \text{ C} = \text{solid}$
Interactions with water	Van der Waals forces	Van der Waals forces	H bonding	H bonding	H bonding (not extensive)	H bonding	H bonding (not extensive)	H bonding	H bonding
Solubility in water	Insoluble	Insoluble	$\leq 4 \text{ C}$ soluble	Limited solubility	Mostly insoluble	$\leq 6 \text{ C}$ soluble	Insoluble	$\leq 5 \text{ C}$ soluble	$\leq 4 \text{ C}$ soluble
Conductivity of aq solution	—	—	Does not conduct	Conducts	—	Conducts	—	Conducts	Does not conduct
Effect on litmus	No change	No change	No change	Blue $\rightarrow$ red	No change	Blue $\rightarrow$ red	No change	Red $\rightarrow$ blue	No change


# H Reagents In Organic Chemistry

Reagent	Conditions	Organic Cps that react	Product (s) formed & observations	Type of Reaction	Equations and Remarks
	uv light (note: $\text{Cl}_2$ also undergoes FRS with alkanes in uv light to form chloroalkane)	Alkane	Bromoalkane Decolourisation of bromine and white fumes of $\text{HBr}$ observed.	Free Radical Substitution	<p>Eg <math>\text{C}_6\text{H}_5\text{CH}_2\text{CH}_3 + \text{Br}_2 \xrightarrow{\text{UV}} \text{C}_6\text{H}_5\text{CHBrCH}_3 + \text{HBr}</math> Any compounds with H attached to a saturated C may undergo FRS. Variety of products formed. To favour monobromination, use excess of the alkane. Major product is the one that is formed via the more stable alkyl radical intermediate. Stability of alkyl radical:</p> $\text{CH}_3\bullet < \text{R}-\dot{\text{C}}\text{H} < \text{R}-\dot{\text{C}}\text{H}-\text{R} < \text{R}-\dot{\text{C}}\text{R}_2$ <p>Stability due to a benzene ring is equivalent to two alkyl groups.</p>
Bromine	$\text{Br}_2$ in $\text{CCl}_4$	Alkene	Dibromoalkane Decolourisation of brown $\text{Br}_2$	Electrophilic addition	<p>Eg: <math>\text{CH}_3\text{CH}=\text{CH}_2 + \text{Br}_2 \rightarrow \text{CH}_3\text{CHBrCH}_2\text{Br}</math> This is used as a distinguishing test for alkenes.</p>
		Phenol	2- or 4-bromophenol Decolourisation of brown $\text{Br}_2$	Easy electrophilic substitution (Electrophilic subst on a highly activated benzene ring in the absence of catalysts)	<p>Eg:</p>  <p>Major product is bromophenol rather than dibromoalkane because <math>\text{H}_2\text{O}</math> is used as the solvent. Addition follows Markovnikov's Rule (may be explained by stability of carbocation.) Eg: <math>\text{CH}_3\text{CH}=\text{CH}_2 + \text{Br}_2 + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{Br} + \text{HBr}</math> Decolourisation of bromine water is used as a distinguishing test for alkenes.</p>
	Aq $\text{Br}_2$	Alkene	Bromohydrin Decolourisation of orange $\text{Br}_2$	Electrophilic addition	
		Phenol or phenylamine	2,4,6-tribromophenol or 2,4,6-tribromophenylamine Decolourisation of orange $\text{Br}_2$ and formation of white ppt.	Easy electrophilic substitution	<p></p> <p><math>\text{X} = \text{OH}</math> or <math>\text{NH}_2</math> Both <math>-\text{OH}</math> and <math>-\text{NH}_2</math> activates the benzene ring towards electrophilic substitution so that the catalyst <math>\text{FeBr}_3</math> is not required. This is used as a distinguishing test for phenol and phenylamine.</p>
		N-phenyl-ethanamide	Decolourisation of orange $\text{Br}_2$	Easy Electrophilic substitution	<p>The <math>-\text{NHCOCH}_3</math> is less activating than the <math>-\text{NH}_2</math> group so that monosubstitution occurs. Hence to convert phenylamine to 2- or 4-bromophenylamine, <math>-\text{NH}_2</math> is first converted to <math>-\text{NHCOCH}_3</math> before bromination. Then the <math>-\text{NHCOCH}_3</math> is hydrolysed back to <math>-\text{NH}_2</math> as shown by the rxn scheme below:</p> 

Reagent	Conditions	Organic Cpd's that react	Product formed & observation	Type of Reaction	Equations and Remarks
Br <sub>2</sub>	FeBr <sub>3</sub> or AlBr <sub>3</sub> as catalyst. (If Cl <sub>2</sub> and FeCl <sub>3</sub> is used, chlorination occurs)	Benzene	Bromobenzene Decolourisation of Br <sub>2</sub> with formation of white fumes.	Electrophilic substitution	Eg  + Br <sub>2</sub> $\xrightarrow{\text{FeBr}_3 \text{ or AlBr}_3}$  + HBr If another group, G, is already present on the benzene ring, substitution is at 2,4 position if G = alkyl group, halogen, -OH, -OR, -NH <sub>2</sub> , -NHR, -NR <sub>2</sub> , -NHCOR and at 3- position if G = -COOH, -COOR, -COR, -NO <sub>2</sub>
Hydrogen halide: HCl or HBr	Anhydrous conditions	Alkene	Halogenoalkane	Electrophilic addition	Addition follows Markovnikov's Rule (may be explained by stability of carbocation). Eg: $\text{CH}_3\text{CH}=\text{CH}_2 + \text{HCl} \rightarrow \text{CH}_3\text{CHClCH}_3$
Conc H <sub>2</sub> SO <sub>4</sub>	xs conc H <sub>2</sub> SO <sub>4</sub> , 170 °C	Alcohol	Alkene	Dehydration (elimination)	Product formed follows Saytzeff's Rule (more substituted alkene is major product because of their greater stability). $\text{xs conc H}_2\text{SO}_4, 170^\circ\text{C} \rightarrow \text{CH}_3\text{CH}=\text{CHCH}_3 + \text{H}_2\text{O}$
	1. Add cold conc H <sub>2</sub> SO <sub>4</sub> 2. Add water and warm	Alkene	Alcohol	Hydration (electrophilic addition followed by nucleophilic substitution)	Addition follows Markovnikov's Rule (may be explained by stability of carbocation). Eg: $\text{CH}_3\text{CH}=\text{CH}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{CH}_3\text{CH}(\text{OSO}_3\text{H})\text{CH}_3$ $\text{CH}_3\text{CH}(\text{OSO}_3\text{H})\text{CH}_3 + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{CH}(\text{OH})\text{CH}_3 + \text{H}_2\text{SO}_4$ Usually, only one equation is used for this two step reaction: $\text{CH}_3\text{CH}=\text{CH}_2 + \text{H}_2\text{O} \xrightarrow[2. \text{H}_2\text{O, warm}]{1. \text{conc H}_2\text{SO}_4} \text{CH}_3\text{CH}(\text{OH})\text{CH}_3$
	Used as a catalyst for esterification. heat	Alcohol + Carboxylic Acid	Ester	Esterification	Eg $\text{RCOOH} + \text{R}'\text{OH} \xrightarrow[\text{Warm/ reflux}]{\text{conc H}_2\text{SO}_4} \text{RCOOR}' + \text{H}_2\text{O}$
Al <sub>2</sub> O <sub>3</sub>	Heat	Alcohol	Alkene	Dehydration, (elimination)	Product formed follows Saytzeff's Rule (more substituted alkene is major product because of their greater stability). $\text{Al}_2\text{O}_3, \text{heat}$ $\text{Eg CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_3 \xrightarrow{\text{Al}_2\text{O}_3, \text{heat}} \text{CH}_3\text{CH}=\text{CHCH}_3 + \text{H}_2\text{O}$
Halogenoalkane, AlX <sub>3</sub> / FeX <sub>3</sub> (not in syllabus)	-	Benzene or substituted benzene	Alkylated benzene	Electrophilic Substitution (Friedel Crafts Alkylation)	If another group, G, is already present on the benzene ring, substitution is at 2,4 position if G = alkyl group, halogen, -OH, -OR, -NH <sub>2</sub> , -NHR, -NR <sub>2</sub> , -NHCOR and at 3- position if G = -COOH, -COOR, -COR, -NO <sub>2</sub> eg  + $\text{CH}_3\text{CHBrCH}_2\text{CH}_3 \xrightarrow{\text{FeBr}_3}$  + HBr


Reagent	Conditions	Organic Cpd that react	Product formed & observation	Type of Reaction	Equations and Remarks
Conc $\text{HNO}_3$	Conc $\text{H}_2\text{SO}_4$ , 55/60 °C for benzene, 30 °C for methylbenzene	Benzene or substituted benzene	Nitrobenzene (appears as a yellow oil with almond smell)	Electrophilic Substitution (Nitration)	<p>If another group, G, is already present on the benzene ring, substitution is at 2,4 position iff G = alkyl group, halogen, -OH, -OR, -NH<sub>2</sub>, -NHR, -NR<sub>2</sub>, -NHCOR and at 3- position if G = -COOH, -COOR, -COR, -NO<sub>2</sub> eg</p>  <p>The -OH group is very activating so that trisubstitution occurs:</p>  <p>If dil <math>\text{HNO}_3</math> is used instead of con nitric acid, mononitration occurs:</p> 
	Room temperature, no catalyst needed	Phenol	2,4,6-trinitrophenol	<b>Easy</b> electrophilic substitution (Nitration)	
$\text{Na}_2\text{CO}_3$ (aq)	room temperature	Carboxylic acid	Carboxylate salt. Effervescence of $\text{CO}_2$ .	Acid-base reaction	$2\text{RCOOH} + \text{CO}_3^{2-} \longrightarrow 2\text{RCOO}^- + \text{CO}_2 + \text{H}_2\text{O}$ <p>This is used as a test for carboxylic acid. Acyl chloride will also give <math>\text{CO}_2</math> but fumes of <math>\text{HCl}</math> is also observed:  <math display="block">\text{RCOCl} + \text{H}_2\text{O} \longrightarrow \text{RCOOH} + \text{HCl}</math> <math display="block">2\text{H}^+ + \text{CO}_3^{2-} \longrightarrow \text{CO}_2 + \text{H}_2\text{O}</math></p> <p>Acid with 6C or more and phenol are insoluble in water. Reaction with <math>\text{NaOH}</math> produces the ionic salt which dissolves in water due to stronger ion-dipole interactions formed between the salt and water:  <math display="block">\text{RCOOH} + \text{OH}^- \longrightarrow \text{RCOO}^- (\text{aq}) + \text{H}_2\text{O}</math> <math display="block">\text{C}_6\text{H}_5\text{OH} + \text{OH}^- \longrightarrow \text{C}_6\text{H}_5\text{O}^- (\text{aq}) + \text{H}_2\text{O}</math> <p>This can be used to distinguish insoluble carboxylic acids or phenol from other compounds.            (NB: phenoxide ion, <math>\text{C}_6\text{H}_5\text{O}^-</math> may also be called phenate)</p> </p>
NaOH (KOH can also be used in place of NaOH)	Aq NaOH, room temperature	Carboxylic acid and phenol	$\text{RCOO}^-$ or phenoxide . If $\text{RCOOH}$ has six or more C atoms, the acid is originally insoluble. On addition of aq NaOH, the insoluble phenol or acid dissolves.	Acid-base reaction	
	Aq NaOH, heat	Halogeno- alkane	Alcohol	Nucleophilic Substitution	$\text{RX} + \text{OH}^- (\text{aq}) \xrightarrow{\text{heat}} \text{ROH} + \text{X}^-$ <p>X directly attached to benzene ring cannot be substituted by <math>\text{OH}^-</math> due to partial double bond character (since lone pair of electrons on X delocalised into benzene ring as p-orbital of X overlaps with <math>\pi</math>-electron cloud of benzene).</p>

Reagent	Conditions	Organic Cpds that react	Product formed & observation	Type of Reaction	Equations and Remarks
NaOH	Aq NaOH, heat	(i) Ester (ii) Amide (iii) Nitrile	Carboxylate salt and (i) alcohol or (ii) NH <sub>3</sub> /amine or (iii) NH <sub>3</sub> . For (ii) and (iii), alkaline gas that turned moist red litmus paper blue was evolved.	Hydrolysis	$\text{RCOOR}' + \text{OH}^- \longrightarrow \text{RCOO}^- + \text{R}'\text{OH}$ $\text{RCONHR}' + \text{OH}^- \longrightarrow \text{RCOO}^- + \text{R}'\text{NH}_2$ $\text{RCN} + \text{OH}^- + \text{H}_2\text{O} \longrightarrow \text{RCOO}^- + \text{NH}_3$ <p>The evolution of ammonia from nitrile and primary amide and the evolution of amines from sec and tert amides (provided these amines are volatile, 5C or less) on warming with aq NaOH may be used as a distinguishing test for amide and nitrile against other classes of compounds.</p>
		Halogeno-alkane	Alkene	Elimination (dehydrohalogenation)	Product formed follows Saytzeff's Rule (more substituted alkene is major product because of their greater stability). Alcoholic NaOH Eg $\text{CH}_3\text{CHBrCH}_2\text{CH}_3 \xrightarrow{\text{Alcoholic NaOH}} \text{CH}_3\text{CH=CHCH}_3 + \text{H}_2\text{O}$
Sodium alkoxide	heat	Halogeno-alkane	Ether	Nucleophilic Substitution	$\text{RX} + \text{R}'\text{O}^- \longrightarrow \text{ROR}' + \text{X}^-$ (solvent used is ROH) Sodium alkoxide is obtained by reacting the appropriate alcohol with Na: $2\text{ROH} + 2\text{Na} \longrightarrow 2\text{R}'\text{O}^-\text{Na}^+ + \text{H}_2$
Conc NH <sub>3</sub>	In ethanol, heat in a sealed tube	Halogeno-alkane	Amine	Nucleophilic Substitution	$\text{RX} + \text{NH}_3 \longrightarrow \text{RNH}_2 + \text{HX}$ Primary amine formed. NH <sub>3</sub> needed to be used in excess to favour formation of 1° amine. With excess RX, R <sub>2</sub> NH, R <sub>3</sub> N or R <sub>4</sub> N <sup>+</sup> X <sup>-</sup> may be formed.
NH <sub>3</sub>	–	Acid halide	Amide	Nucleophilic Substitution	$\text{RCOX} + \text{NH}_3 \longrightarrow \text{RCONH}_2 + \text{HX}$ Primary amide formed.
KCN (or NaCN)	Aqueous alcoholic KCN, heat	Halogeno-alkane	Nitrile	Nucleophilic Substitution	$\text{RX} + \text{CN}^- \longrightarrow \text{RCN} + \text{X}^-$ Alcohol is used as a solvent to dissolve both CN <sup>-</sup> and RX. The resultant compound has one C more than the starting RX. This is an important synthesis step allowing length of C chain to be extended by 1. RCN can be hydrolysed to RCOOH (warming RCN with dil acid) or reduce to RCH <sub>2</sub> NH <sub>2</sub> (using LiAlH <sub>4</sub> ). HCN cannot be used as it is not strong enough a nucleophile to react with RX.
HCN	Trace amt of KCN or small amount of aq NaOH, cold	Carbonyl compound (ie aldehyde or ketone)	Cyanohydrin	Nucleophilic addition	$\text{R}-\overset{\text{O}}{\underset{\text{  }}{\text{C}}}-\text{R}' + \text{HCN} \xrightarrow{\text{trace KCN}} \text{R}-\overset{\text{OH}}{\underset{\text{  }}{\text{C}}}-\text{R}' \quad \text{R}, \text{R}' = \text{H, alkyl or aryl}$ <p>Cyanohydrin may be hydrolysed (by warming with dil acid) to give 2-hydroxycarboxylic acid or reduced to 2-hydroxyamine. This reaction allows an increase of one carbon in carbon chain length.</p>
Dilute acid (HCl, HNO <sub>3</sub> or H <sub>2</sub> SO <sub>4</sub> )	heat acid with compound	(i) Ester (ii) Amide (iii) Nitrile	Carboxylic acid and (i) alcohol or (ii) NH <sub>4</sub> <sup>+</sup> /amine salt or (iii) NH <sub>4</sub> <sup>+</sup>	Hydrolysis	$\text{RCOOR}' + \text{H}_2\text{O} \longrightarrow \text{RCOOH} + \text{R}'\text{OH}$ $\text{RCONHR}' + \text{H}_2\text{O} + \text{H}^+ \longrightarrow \text{RCOOH} + \text{R}'\text{NH}_3^+$ $\text{RCN} + \text{H}_2\text{O} + \text{H}^+ \longrightarrow \text{RCOOH} + \text{NH}_4^+$

Reagent	Conditions	Type of Organic Cps that react	Product formed & observation	Type of Reaction	Equations and Remarks
RCOCl (RCOBr may also be used)	Room temperature	(i) alcohol (ii) phenol (iii) amine (either pri or sec)	(i) ester (ii) phenolic ester (iii) amide	(i) and (ii) esterification; (iii) amide formation (condensation)	<p>(i) <math>\text{RCOCl} + \text{R}'\text{OH} \longrightarrow \text{RCOOR}' + \text{HCl}</math></p> <p>(ii) <math>\text{RCOCl} + \text{C}_6\text{H}_5\text{OH} \longrightarrow \text{RCOOC}_6\text{H}_5 + \text{HCl}</math></p> <p>(iii) <math>\text{RCOCl} + \text{R}'\text{NH}_2 \longrightarrow \text{RCONHR}' + \text{HCl}</math></p> <p>To convert phenol to phenolic ester, acid chloride must be used. Carboxylic acids are not reactive enough to react with phenol. To convert amine to amide, acid chloride is needed. Carboxylic acid does not react with amine to form amide (unless under industrial conditions) but merely neutralises it.</p>
Aq $\text{KMnO}_4$	Cold dil alkaline	Alkene	Diol. Formation of black $\text{MnO}_2$ .		<p>Eg <math>\text{CH}_3\text{CH}=\text{CH}_2 + \text{H}_2\text{O} + [\text{O}] \longrightarrow \text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{OH}</math></p> <p>C=C bond cleaves. Final product dependent on groups attached to each C in the C=C bond: <math>=\text{CH}_2 \rightarrow \text{CO}_2 + \text{H}_2\text{O}</math>; <math>=\text{CHR} \rightarrow \text{RCOOH}</math>; <math>=\text{CRR}' \rightarrow \text{RCOR}'</math>.</p>
		Alkene (C=C)	$\text{CO}_2$ , carboxylic acid or ketone. Decolourisation of purple $\text{KMnO}_4$		<p>Eg </p>
		Alkyl benzene	Benzoic acid and $\text{CO}_2$		<p>Alkylbenzene will be oxidized to benzoic acid, eg  <math>\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{CH}_3 + 9[\text{O}] \longrightarrow \text{C}_6\text{H}_5\text{COOH} + 2\text{CO}_2 + 3\text{H}_2\text{O}</math>            The alkyl group needed to have a H atom on C attached to benzene ring for alkylbenzene to be oxidised. Hence, <math>\text{C}(\text{CH}_3)_3\text{C}_6\text{H}_5</math> would not be oxidised.            These are the only two common carboxylic acids which can be oxidized by <math>\text{KMnO}_4</math>: <math>\text{HCOOH} + [\text{O}] \longrightarrow \text{CO}_2 + \text{H}_2\text{O}</math>  <math>\text{HOOC-COOH} + [\text{O}] \longrightarrow 2\text{CO}_2 + \text{H}_2\text{O}</math></p>
		$\text{HCOOH}$ , and ethanedioic acid,	$\text{CO}_2$ and $\text{H}_2\text{O}$		<p><math>\text{RCHO} + [\text{O}] \longrightarrow \text{RCOOH}</math>  <math>\text{RCH}_2\text{OH} + 2[\text{O}] \longrightarrow \text{RCOOH} + \text{H}_2\text{O}</math>  <math>\text{RCH}(\text{OH})\text{R}' + [\text{O}] \longrightarrow \text{RCR}'=\text{O} + \text{H}_2\text{O}</math></p>
		aldehyde, $1^\circ$ or $2^\circ$ alcohol	Carboxylic acid (from aldehyde & $1^\circ$ ROH) & ketone (fr $2^\circ$ ROH)		<p>heat with dil <math>\text{H}_2\text{SO}_4</math>, hydrolyses the ester into an acid and alcohol, if either the acid or alcohol can undergo oxidation (acid = <math>\text{HCOOH}</math> or ethanedioic acid; alcohol = <math>\text{CH}_3\text{OH}</math>, pri or sec alcohol), decolourisation of <math>\text{KMnO}_4</math> observed.            Eg (1) <math>\text{HCOOC}(\text{CH}_3)_3 + \text{H}_2\text{O} \longrightarrow \text{HCOOH} + \text{C}(\text{CH}_3)_3\text{OH}</math>  <math>\text{HCOOH} + [\text{O}] \longrightarrow \text{CO}_2 + \text{H}_2\text{O}</math>            (2) <math>\text{CH}_3\text{COOCH}_2\text{CH}_2\text{CH}_3 + \text{H}_2\text{O} \longrightarrow \text{CH}_3\text{COOH} + \text{CH}_3\text{CH}_2\text{CH}_2\text{OH}</math>  <math>\text{CH}_3\text{CH}_2\text{CH}_2\text{OH} + 2[\text{O}] \longrightarrow \text{CH}_3\text{CH}_2\text{COOH} + \text{H}_2\text{O}</math></p>
Add dilute $\text{H}_2\text{SO}_4$ and $\text{KMnO}_4$ to the compound and heat		Esters which hydrolyses to form $\text{HCOOH}$ , ethanedioic acid, $\text{CH}_3\text{OH}$ , $1^\circ$ or $2^\circ$ alcohol	$\text{CO}_2$ (from $\text{HCOOH}$ , ethanedioic acid or $\text{CH}_3\text{OH}$ ), carboxylic acid (from $1^\circ$ alcohol) or ketone (from $2^\circ$ alcohol)		
		Phenylketone and $3^\circ$ ROH with phenyl gp attached to hydroxyl C.	Benzoic acid		<p>Phenyl ketone and <math>3^\circ</math> alcohol with phenyl gp attached to hydroxyl C contain a benzene ring and can be oxidised to benzoic acid by heating with <math>\text{KMnO}_4/\text{H}^+</math>:            eg <math>\text{C}_6\text{H}_5\text{COCH}_2\text{CH}_3 + 7[\text{O}] \longrightarrow \text{C}_6\text{H}_5\text{COOH} + 2\text{CO}_2 + 2\text{H}_2\text{O}</math>  <math>\text{C}_6\text{H}_5\text{C}(\text{OH})(\text{CH}_3)_2 + 8[\text{O}] \longrightarrow \text{C}_6\text{H}_5\text{COOH} + 2\text{CO}_2 + 3\text{H}_2\text{O}</math></p>


Reagent	Conditions	Type of Organic Cpd's that react	Product formed & observation	Type of Reaction	Equations and Remarks
Aq $K_2Cr_2O_7$	Add dil $H_2SO_4$ and aq $K_2Cr_2O_7$ to compound and heat	$HCOOH$ , aldehyde, $1^\circ$ & $2^\circ$ alcohols, and esters which form oxidisable products on hydrolysis.	Products same as $KMnO_4$ . Observation: Colour change from orange to green.	Oxidation	Potassium dichromate(VI) is a weaker oxidizing agent than $KMnO_4$ and is usually not used to oxidize alkenes, alkylbenzenes and ethanedioic acid.
	Add dil $H_2SO_4$ and aq $K_2Cr_2O_7$ to compound. Immediate distillation	Primary alcohol	Aldehyde	Oxidation	$RCH_2OH + [O] \longrightarrow RCHO + H_2O$
$H_2$	Ni, heat	Alkene	Alkane	Addition/reduction	$RCH=CHR' + H_2 \longrightarrow RCH_2CH_2R'$
		Nitrile, RCN	Primary amine	reduction	$RCN + 2H_2 \longrightarrow RCH_2NH_2$
		Aldehyde and ketone	Primary and secondary alcohol	Reduction	$RCHO + H_2 \longrightarrow RCH_2OH$ $RCR'O + H_2 \longrightarrow RCH(OH)R'$ (does not reduce $RCOOH$ !)
		(i) Nitrile, (ii) aldehyde, (iii) ketone and (iv) carboxylic acid	(i) Primary amine, (ii) primary alcohol, (iii) secondary alcohol and (iv) primary alcohol.	Reduction	$RCN + 4[H] \longrightarrow RCH_2NH_2$ $RCHO + 2[H] \longrightarrow RCH_2OH$ $RCR'O + 2[H] \longrightarrow RCH(OH)R'$ $RCOOH + 4[H] \longrightarrow RCH_2OH + H_2O$
$LiAlH_4$ in dry ether	Room temp	aldehyde, ketone	primary alcohol, secondary alcohol	Reduction	$RCHO + 2[H] \longrightarrow RCH_2OH$ $RCR'O + 2[H] \longrightarrow RCH(OH)R'$
$NaBH_4$ in methanol	Room temp	Nitrobenzene	Phenylamine	Reduction	$C_6H_5NO_2 + 6[H] \longrightarrow C_6H_5NH_2 + 2H_2O$ (aq NaOH is used in $2^{nd}$ step to liberate free amine from amine salt, $C_6H_5NH_3^+ Cl^-$ )
Sn and conc HCl	Reflux	Alcohol	Chloroalkane	N. Substitution	$ROH + PCl_5 \longrightarrow RCl + HCl + POCl_3$
	anhydrous	Carboxylic acid	Acyl chloride	N. Substitution	$RCOOH + PCl_5 \longrightarrow RCOCl + HCl + POCl_3$ $PCl_5$ replaces OH group in ROH or $RCOOH$ with Cl
$PCl_3$ or $PBr_3$	anhydrous	Alcohol	Chloroalkane/bromoalkane	N. Substitution	$3ROH + PX_3 \longrightarrow 3RX + H_3PO_3$ where X = Br or Cl
	anhydrous	Carboxylic acid	Acid chloride/bromide	N. Substitution	$3RCOOH + PX_3 \longrightarrow 3RCOX + H_3PO_3$ where X = Br or Cl $PX_3$ replaces OH group in ROH or $RCOOH$ with X where X = Br or Cl
$SOCl_2$	anhydrous	Alcohol	Chloroalkane	N. substitution	$ROH + SOCl_2 \longrightarrow RCl + HCl + SO_2$ $SOCl_2$ replaces OH group in ROH with Cl

# I Reagents used to Test for Various Functional Groups

Reagent	Type of Organic Cpds that react	Test and Conditions	Observations	Equation and Remarks
aq $\text{KMnO}_4$ + dil $\text{H}_2\text{SO}_4$ (Note: This test is not specific and more specific tests are usually preferred)	Alkene ( $\text{C}=\text{C}$ )	Heat compound with acidified $\text{KMnO}_4$ (aq) (dil $\text{H}_2\text{SO}_4$ is used for acidification)	Purple $\text{KMnO}_4$ was decolourised.	Eg: $\text{CH}_3\text{CH}=\text{CH}_2 + 5[\text{O}] \rightarrow \text{CH}_3\text{COOH} + \text{CO}_2 + \text{H}_2\text{O}$ Eg: $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_3 + 6[\text{O}] \rightarrow \text{C}_6\text{H}_5\text{COOH} + \text{CO}_2 + 2\text{H}_2\text{O}$ $\text{HCOOH}$ and ethanedioic acid are oxidised to $\text{CO}_2$ and $\text{H}_2\text{O}$ . Aldehydes and primary alcohols are oxidised to carboxylic acids while secondary alcohols are oxidised to ketones. Eg $(\text{CH}_3)_2\text{CHOH} + [\text{O}] \rightarrow (\text{CH}_3)_2\text{C}=\text{O} + \text{H}_2\text{O}$
	Alkyl benzene			Ester hydrolyses to form acid and alcohol on being warmed with dil $\text{H}_2\text{SO}_4$ . If either the acid or alcohol can be oxidized, $\text{KMnO}_4$ will be decolourised. Eg $\text{CH}_3\text{COOCH}_3 + \text{H}_2\text{O} + \text{H}^+ \rightarrow \text{CH}_3\text{COOH} + \text{CH}_3\text{OH}$ $\text{CH}_3\text{OH} + 3[\text{O}] \rightarrow \text{CO}_2 + 2\text{H}_2\text{O}$
	HCOOH, ethanedioic acid, aldehyde, primary or secondary alcohol			Eg: $\text{CH}_3\text{CH}=\text{CH}_2 + \text{Br}_2 \rightarrow \text{CH}_3\text{CH}_2\text{BrCH}_2\text{Br}$ Addition follows Markovnikov's Rule Eg: $\text{CH}_3\text{CH}=\text{CH}_2 + \text{Br}_2 + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{Br} + \text{HBr}$
Bromine	Alkene ( $\text{C}=\text{C}$ )	React compound with $\text{Br}_2$ in $\text{CCl}_4$	Brown $\text{Br}_2$ was rapidly decolourised.	 <p>where <math>\text{X} = \text{OH}</math> or <math>\text{NH}_2</math></p> <p>Alkenes, phenol and phenylamine will also decolourise bromine. This test is used to distinguish compound with benzene rings from those that do not have benzene ring and <math>\text{C}=\text{C}</math> bond, eg benzene from cyclohexane.</p> <p>Eg: <math>2\text{CH}_3\text{COOH} + \text{CO}_3^{2-} \rightarrow 2\text{CH}_3\text{COO}^- + \text{H}_2\text{O} + \text{CO}_2</math> Acyl chloride will also give effervescence of <math>\text{CO}_2</math> too but fumes of <math>\text{HCl}</math> is also evolved: Eg: <math>\text{CH}_3\text{COCl} + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{COOH} + \text{HCl}</math> <math>2\text{H}^+ + \text{CO}_3^{2-} \rightarrow \text{CO}_2 + \text{H}_2\text{O}</math></p>
	Alkene ( $\text{C}=\text{C}$ )	React compound with aqueous bromine	Orange / brown $\text{Br}_2$ was rapidly decolourised, white fumes of $\text{HBr}$ and white ppt.	
	Phenol and phenylamine	React compound with bromine and $\text{FeCl}_3$ as catalyst in the dark	Brown $\text{Br}_2$ was decolourised.	
Aqueous $\text{Na}_2\text{CO}_3$	Carboxylic acid	Add aq $\text{Na}_2\text{CO}_3$ to compound	Effervescence of $\text{CO}_2$ [gave white ppt with aq $\text{Ca}(\text{OH})_2$ ].	










Reagent	Type of Organic Cpds that react	Test and Conditions	Observations	Equation and Remarks
Aqueous NaOH	Phenol	Add aq NaOH to compound	Insoluble phenol dissolves to form a colourless soln (1 miscible layer).	Phenol is sparingly soluble but on conversion to sodium phenoxide by reaction with aq NaOH, will dissolve: $C_6H_5OH(s) + NaOH(aq) \rightarrow C_6H_5O^-Na^+(aq) + H_2O(l)$ This test can be used to distinguish phenol from solids that are insoluble in aq NaOH but will not be able to distinguish phenol from benzoic acid.
	Amine salt	Add aq NaOH to a solution of the compound. Warm gently.	A gas that turned moist red litmus paper as evolved (for volatile amine - with 5 C or less eg $CH_3NH_2$ ) or an organic layer separates out (non-volatile amines such as phenylamine)	Eg: $CH_3NH_3^+(aq) + OH^-(aq) \rightarrow CH_3NH_2(g) + H_2O(l)$ Eg: $C_6H_5NH_3^+(aq) + OH^-(aq) \rightarrow C_6H_5NH_2(l) + H_2O(l)$
	Ammonium salt	Add aq NaOH to a solution of the compound. Warm gently.	A gas that turns moist red litmus paper blue was evolved.	Ammonia was formed. The gentle warming was necessary to drive the soluble $NH_3$ out of solution. Eg: $NH_4^+(aq) + OH^-(aq) \rightarrow NH_3(g) + H_2O(l)$
	Amide	Add aq NaOH to the compound. Heat strongly	After some time of warming, a gas that turns moist red litmus paper blue was evolved.	Only amides that hydrolyse to give volatile ammonia or amines (5C or less) will give a gas that turns moist red litmus blue. The strong heating is needed to hydrolyse the amide bond: Eg $CH_3CON(CH_3)_2 + OH^- \rightarrow CH_3COO^- + NH(CH_3)_2 \uparrow$
Aqueous NaOH (with dil $HNO_3$ & aq $AgNO_3$ )	Halogenoalkane	Heat cpd with aq NaOH. Cool. Acidify with dil $HNO_3$ . Add aq $AgNO_3$ .	White ppt of $AgCl$ (for $RCI$ ) Cream ppt of $AgBr$ (for $RBr$ ) Yellow ppt for $AgI$ (for $RI$ )	Eg: $C_6H_5CH_2Cl + OH^- \rightarrow C_6H_5CH_2OH + Cl^-$ , $Ag^+(aq) + Cl^-(aq) \rightarrow AgCl(s)$ Test is negative if $Cl$ , $Br$ or $I$ is directly attached to benzene ring.
	Acid halide	Add aq $AgNO_3$ to compound at r.t.	White fumes seen with formation of ppt: White ppt of $AgCl$ (for $RCOCl$ ) Cream ppt of $AgBr$ (for $RCOBr$ ) Yellow ppt for $AgI$ (for $RCOI$ )	Eg: $RCOCl + H_2O \rightarrow RCOOH(aq) + HCl(aq)$ $Ag^+(aq) + Cl^-(aq) \rightarrow AgCl(s)$
Sodium	Alcohol	Add sodium to compound	Effervescence of $H_2$ (gas extinguished lighted splinter with a pop sound)	Eg: $C_2H_5OH + Na \rightarrow C_2H_5O^-Na^+ + \frac{1}{2}H_2(g)$ Note that Na should not be used unless there is no other alternative as it reacts explosively with water, phenol and carboxylic acid. A better test for alcohol with a $CH_3CH(OH)-$ group is the $CHI_3$ test (warming with aq alkaline $I_2$ ).

Reagent	Type of Organic Cpd's that react	Test and Conditions	Observations	Equation and Remarks
Iodine solution + aq NaOH	Alcohol with $\text{CH}_3\text{C}(\text{OH})\text{H}$ —gp or carbonyl cpd with $\text{CH}_3\text{CO}$ —gp	Warm compound with aqueous iodine and aqueous NaOH	Yellow ppt of $\text{CHI}_3$ formed.	$\text{RCH}(\text{OH})\text{CH}_3 + 4\text{I}_2 + 6\text{OH}^- \rightarrow \text{RCOO}^- + \text{CHI}_3(\text{s}) + 5\text{I}^- + 5\text{H}_2\text{O}$ $\text{RCOCH}_3 + 3\text{I}_2 + 4\text{OH}^- \rightarrow \text{RCOO}^- + \text{CHI}_3(\text{s}) + 3\text{I}^- + 3\text{H}_2\text{O}$ The ppt is a carboxylate salt with one C less than the starting alcohol (ethanol of a secondary 2-ol) or carbonyl cpd (ethanal or methylketone). Note that methyl esters or amides <b>DO NOT</b> give $\text{CHI}_3$ with $\text{I}_2/\text{OH}^-$ .
Neutral $\text{FeCl}_3$ (aq)	Phenol	Add neutral $\text{FeCl}_3$ (aq) to the compound	Violet complex formed.	$6\text{C}_6\text{H}_5\text{OH} + \text{Fe}^{3+} \rightarrow [\text{Fe}(\text{C}_6\text{H}_5\text{O})_6]^{3-} + 6\text{H}^+$ Eg: 
2,4-dinitrophenylhydrazine (2,4-DNPH or Brady's reagent)	Carbonyl cpds (aldehyde or ketone)	Add 2,4-dinitrophenylhydrazine to compound	Orange ppt formed.	Note: esters, amides, carboxylic acids <b>give NO ppt</b> with 2,4-dinitrophenylhydrazine even though they contain $\text{C}=\text{O}$ group.
Fehling's solution	Aliphatic aldehyde	Warm cpd with Fehling's solution.	Reddish brown ppt of $\text{Cu}_2\text{O}$ formed.	$\text{RCHO} + 2\text{Cu}^{2+} + 5\text{OH}^- \rightarrow \text{RCOO}^- + \text{Cu}_2\text{O} + 3\text{H}_2\text{O}$ A carboxylate salt with the same number of C as the aldehyde is formed. Methanoic acid (with the $\text{H}-\text{C}=\text{O}$ group) also reacts with Fehling's solution (and Tollens' reagent), itself being oxidised to $\text{CO}_3^{2-}$ . $\text{HCOOH} + 2\text{Cu}^{2+} + 6\text{OH}^- \rightarrow \text{CO}_3^{2-} + \text{Cu}_2\text{O} + 4\text{H}_2\text{O}$ Benzaldehyde (aromatic aldehydes) <b>DOES NOT</b> react with Fehling's solution.
Tollens' reagent	Aldehyde	Warm cpd with Tollens' solution.	Silver mirror formed.	$\text{RCHO} + 2\text{Ag}(\text{NH}_3)_2^+ + 3\text{OH}^- \rightarrow \text{RCOO}^- + 2\text{Ag} + 4\text{NH}_3 + 2\text{H}_2\text{O}$ A carboxylate salt with the same no of C as the aldehyde is formed. Benzaldehyde reacts with Tollens' reagent with much less ease and a grey ppt may be obtained instead of a silver mirror. Methanoic acid is oxidised to carbonate ion: $\text{HCOOH} + 2\text{Ag}(\text{NH}_3)_2^+ + 4\text{OH}^- \rightarrow \text{CO}_3^{2-} + 2\text{Ag} + 4\text{NH}_3 + 3\text{H}_2\text{O}$

## J SUMMARY OF COMMON REAGENTS

Table 1: Common reagents used for identifying various functional groups

Organic Compound	Alkene	Alkyl benzene	Halogeno alkane	Alcohol		Phenol	Carbonyl compound				Carboxylic acid RCOOH
				1°/2° ROH	3° ROH		Aldehyde RCHO	Benzaldehyde C <sub>6</sub> H <sub>5</sub> CHO	Ketone RCOR	Carbonyl with CH <sub>3</sub> CO- structure	
Functional Group	C=C		RX	RCH <sub>2</sub> OH or R <sub>2</sub> CHOH	R <sub>3</sub> COH						
Example Br <sub>2</sub> (aq)	CH <sub>2</sub> =CH <sub>2</sub> decolourise		CH <sub>3</sub> CH <sub>2</sub> Br	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> OH	(CH <sub>3</sub> ) <sub>3</sub> COH	Decolourise, white ppt Violet colouration	CH <sub>3</sub> CH <sub>2</sub> CHO			CH <sub>3</sub> COCH <sub>3</sub>	CH <sub>3</sub> COOH
FeCl <sub>3</sub> neutral											
KMnO <sub>4</sub> /H <sub>2</sub> SO <sub>4</sub> , heat	Purple MnO <sub>4</sub> <sup>-</sup> decolourise	Purple MnO <sub>4</sub> <sup>-</sup> decolourise		Purple MnO <sub>4</sub> <sup>-</sup> decolourise			Purple MnO <sub>4</sub> <sup>-</sup> decolourise	Purple MnO <sub>4</sub> <sup>-</sup> decolourise			
K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> /H <sub>2</sub> SO <sub>4</sub> , heat				Orange → green			Orange → green	Orange → green			
Na				H <sub>2</sub> evolved	H <sub>2</sub> evolved	H <sub>2</sub> evolved					H <sub>2</sub> evolved
NaOH(aq)						Phenol dissolves					Heat evolved
Na <sub>2</sub> CO <sub>3</sub> (aq)											CO <sub>2</sub> evolved
I <sub>2</sub> (aq)/NaOH(aq), heat										Yellow ppt (iodoform)	
2,4-DNPH							Orange ppt	Orange ppt	Orange ppt	Orange ppt	
Tollens' reagent, warm						Silver mirror	Silver mirror	Silver mirror			
Fehling's solution, warm							Reddish brown ppt				
NaOH(aq), heat, cool, acidify, AgNO <sub>3</sub> (aq)			White ppt ⇒ X = Cl, Cream ppt ⇒ X = Br, Yellow ppt ⇒ X = I								

R = alkyl group

- Test for RCOX: Add water, white fumes of HX formed. OR Add aqueous AgNO<sub>3</sub>. White fumes and white, cream or yellow ppt for X = Cl, Br or I respectively.
- Test for salts of amine, eg RNH<sub>3</sub><sup>+</sup>NO<sub>3</sub><sup>-</sup>. Heat gently with aq NaOH. If R contains 5C or less, gas evolved turned moist red litmus paper blue. For R with 6C or more, a immiscible layer formed.
- Test for amide: RCONH<sub>2</sub> Heat strongly with aq NaOH. Gas evolved turned moist red litmus paper blue.

Table 2: Reactions with common oxidising agents



	KMnO <sub>4</sub> (aq), H <sub>2</sub> SO <sub>4</sub> (aq), heat	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> (aq), H <sub>2</sub> SO <sub>4</sub> (aq), heat	KMnO <sub>4</sub> (aq), NaOH(aq), heat	KMnO <sub>4</sub> (aq), NaOH(aq), Cold	Remarks/Equations
<b>Alkene</b>	✓ Purple KMnO <sub>4</sub> decolourised	✗	✓ Purple KMnO <sub>4</sub> decolourised, brown MnO <sub>2</sub> ppt	✓ Purple KMnO <sub>4</sub> decolourised; brown MnO <sub>2</sub> ppt	CO <sub>2</sub> may be produced for terminal alkenes E.g. CH <sub>3</sub> CH=CH <sub>2</sub> + 5[O] → CH <sub>3</sub> COOH + CO <sub>2</sub> + H <sub>2</sub> O. Oxidative cleavage of alkenes with hot KMnO <sub>4</sub> .
<b>Alkybenzene</b>	✓ Purple KMnO <sub>4</sub> decolourised	✗	✓ Purple KMnO <sub>4</sub> decolourised; benzoate salt formed	✗	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> CH <sub>3</sub> + 6[O] → C <sub>6</sub> H <sub>5</sub> COOH + CO <sub>2</sub> + 2H <sub>2</sub> O Oxidation only occurs when alkyl side-gp has H atom on C attached to benzene ring, i.e.  and 
<b>1°/2° Alcohol</b>	✓ Purple KMnO <sub>4</sub> decolourised	✓ Orange K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> turns green	✓ Purple KMnO <sub>4</sub> decolourised	✗	1° alc are oxidised to carboxylic acids 1° alc are oxidised to aldehydes using K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> /H <sup>+</sup> and heat with immediate distillation. 2° alc are oxidised to ketones. E.g. (CH <sub>3</sub> ) <sub>2</sub> CHOH + [O] → (CH <sub>3</sub> ) <sub>2</sub> C=O + H <sub>2</sub> O
<b>3° Alcohol</b>	✗	✗	✗	✗	3° alcohols are generally resistant to oxidation except for 3° alcohols with phenyl gp attached to hydroxyl C contain a benzene ring and can be oxidised to benzoic acid by warming with KMnO <sub>4</sub> /H <sup>+</sup> . E.g. C <sub>6</sub> H <sub>5</sub> C(OH)(CH <sub>3</sub> ) <sub>2</sub> + 8[O] → C <sub>6</sub> H <sub>5</sub> COOH + 2CO <sub>2</sub> + 3H <sub>2</sub> O
<b>Aldehyde</b>	✓ Purple KMnO <sub>4</sub> decolourised	✓ Orange K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> turns green	✓ Purple KMnO <sub>4</sub> decolourised; salt formed	✗	Aldehydes are oxidised to carboxylic acids E.g. CH <sub>3</sub> CHO + [O] → CH <sub>3</sub> COOH
<b>Benzaldehyde</b>	✓ Purple KMnO <sub>4</sub> decolourised	✓ Orange K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> turns green	✓ Purple KMnO <sub>4</sub> decolourised; benzoate salt formed	✗	E.g. C <sub>6</sub> H <sub>5</sub> CHO + [O] → C <sub>6</sub> H <sub>5</sub> COOH
<b>Ketone</b>	✗ Purple KMnO <sub>4</sub> decolourised for phenyl ketones	✗	✗	✗	Most ketones are resistant to oxidation except for phenyl ketones which can be oxidised to benzoic acid by warming with KMnO <sub>4</sub> /H <sup>+</sup> . E.g. C <sub>6</sub> H <sub>5</sub> COCH <sub>2</sub> CH <sub>3</sub> + 7[O] → C <sub>6</sub> H <sub>5</sub> COOH + 2CO <sub>2</sub> + 2H <sub>2</sub> O
<b>Ester</b>	✓ Purple KMnO <sub>4</sub> decolourised;	✗	✓ Purple KMnO <sub>4</sub> decolourised	✗	Warming with dil H <sub>2</sub> SO <sub>4</sub> hydrolyses the ester into an acid and alcohol. If either the acid or alcohol can undergo oxidation (acid = HCOOH or ethanedioic acid; alcohol = CH <sub>3</sub> OH, 1° or 2° alcohol), decolourisation of KMnO <sub>4</sub> observed. E.g. (1) HCOOC(CH <sub>3</sub> ) <sub>3</sub> + H <sub>2</sub> O → HCOOH + C(CH <sub>3</sub> ) <sub>3</sub> OH HCOOH + [O] → CO <sub>2</sub> + H <sub>2</sub> O (2) CH <sub>3</sub> COOCH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub> + H <sub>2</sub> O → CH <sub>3</sub> COOH + CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> OH CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> OH + 2[O] → CH <sub>3</sub> CH <sub>2</sub> COOH + H <sub>2</sub> O

Table 3: Reactions with common reducing agents

	H <sub>2</sub> , Ni, heat	LiAlH <sub>4</sub> , dry ether	NaBH <sub>4</sub> in methanol	Sn/conc HCl	Remarks/Equations
Alkene	✓	✗	✗	✗	Reaction can be considered as catalytic hydrogenation/addition/reduction (but <b>NOT</b> electrophilic addition). E.g. $\text{RCH=CHR}' + \text{H}_2 \longrightarrow \text{RCH}_2\text{CH}_2\text{R}'$
Benzene	✓ *	✗	✗	✗	*Benzene will be reduced to cyclohexane only when heated with H <sub>2</sub> Ni under high temperature and pressure (100–200°C, 1500 psi) $\text{C}_6\text{H}_6 + 3\text{H}_2 \longrightarrow \text{C}_6\text{H}_{12}$
Nitrobenzene	✗	✗	✗	✓	E.g. $\text{C}_6\text{H}_5\text{NO}_2 + 6[\text{H}] \longrightarrow \text{C}_6\text{H}_5\text{NH}_2 + 2\text{H}_2\text{O}$ phenylamine formed NaOH(aq) is used to liberate free amine (C <sub>6</sub> H <sub>5</sub> NH <sub>2</sub> ) from amine salt (C <sub>6</sub> H <sub>5</sub> NH <sub>3</sub> <sup>+</sup> Cl <sup>-</sup> ) formed.
Nitrile	✓	✓	✗	✗	Reduction of nitrile always leads to formation of 1° amine. H <sub>2</sub> /Ni, heat: $\text{RCN} + \text{H}_2 \longrightarrow \text{RCH}_2\text{NH}_2$ LiAlH <sub>4</sub> , dry ether: $\text{RCHO} + 2[\text{H}] \longrightarrow \text{RCH}_2\text{OH}$
Aldehyde	✓	✓	✓	✗	H <sub>2</sub> /Ni, heat: $\text{RCHO} + \text{H}_2 \longrightarrow \text{RCH}_2\text{OH}$ LiAlH <sub>4</sub> , dry ether or NaBH <sub>4</sub> $\text{RCHO} + 2[\text{H}] \longrightarrow \text{RCH}_2\text{OH}$
Ketone	✓	✓	✓	✗	H <sub>2</sub> /Ni, heat: $\text{RCOR}' + \text{H}_2 \longrightarrow \text{RCH(OH)R}'$ LiAlH <sub>4</sub> , dry ether or NaBH <sub>4</sub> $\text{RCOR}' + 2[\text{H}] \longrightarrow \text{RCH(OH)R}'$
Carboxylic Acid	✗	✓	✗	✗	Carboxylic acid cannot be reduced by H <sub>2</sub> / Ni. Need to use LiAlH <sub>4</sub> . $\text{RCOOH} + 4[\text{H}] \longrightarrow \text{RCH}_2\text{OH} + \text{H}_2$

**Table 4: Common reagents used to identify some acidic compounds**

	Na(s)	NaOH(aq)	Na <sub>2</sub> CO <sub>3</sub> (aq) or NaHCO <sub>3</sub> (aq)	PCl <sub>5</sub>	Remarks/Equations
Alcohol	✓ H <sub>2</sub> evolves	✗	✗	✓ white fumes of HCl	PCl <sub>5</sub> cannot distinguish between alcohol and carboxylic acid.  ROH + PCl <sub>5</sub> → RCl + POCl <sub>3</sub> + HCl
Phenol	✓ H <sub>2</sub> evolves	✓ Phenol dissolves	✗	✗	C <sub>6</sub> H <sub>5</sub> OH (s) + NaOH (aq) → C <sub>6</sub> H <sub>5</sub> O <sup>-</sup> Na <sup>+</sup> (aq) + H <sub>2</sub> O (l)  This test can be used to distinguish phenol from solids that are insoluble in aq NaOH but will not be able to distinguish phenol from benzoic acid.
Carboxylic Acid	✓ H <sub>2</sub> evolves	✓	✓ CO <sub>2</sub> evolves	✓ white fumes of HCl	E.g. 2CH <sub>3</sub> COOH + CO <sub>3</sub> <sup>2-</sup> → 2CH <sub>3</sub> COO <sup>-</sup> + H <sub>2</sub> O + CO <sub>2</sub>  Acid chloride will also give effervescence of CO <sub>2</sub> with aq Na <sub>2</sub> CO <sub>3</sub> too but fumes of HCl is also evolved: E.g. CH <sub>3</sub> COCl + H <sub>2</sub> O + → CH <sub>3</sub> COOH + HCl 2H <sup>+</sup> + CO <sub>3</sub> <sup>2-</sup> → CO <sub>2</sub> + H <sub>2</sub> O

Table 5: Common reagents used to identify carbonyl compounds

	2,4-DNPH	Tollens' reagent, warm [Ag(NH <sub>3</sub> ) <sub>2</sub> ] <sup>+</sup> , OH <sup>-</sup>	Fehling's reagent, warm Cu <sup>2+</sup> , OH <sup>-</sup>	I <sub>2</sub> (aq), NaOH(aq), warm	Remarks/Equations
<b>Aldehyde (aliphatic)</b>	✓ Orange ppt	✓ Ag mirror	✓ Reddish brown ppt of Cu <sub>2</sub> O	✓ (see remarks) Yellow ppt of CHI <sub>3</sub> only for CH <sub>3</sub> CHO	Ethanal is the only aldehyde that gives yellow ppt of CHI <sub>3</sub> when heated with aq. alkaline I <sub>2</sub> .
<b>Benzaldehyde (aromatic aldehydes : CHO gp bonded directly to benzene ring)</b>	✓ Orange ppt	✓ Ag mirror	✗	✗	Benzaldehyde <b><u>DOES NOT</u></b> react with Fehling's solution.
<b>Ketone</b>	✓ Orange ppt	✗	✗	✓ (see remarks) Yellow ppt of CHI <sub>3</sub> only for methyl ketones (CH <sub>3</sub> CO-)	Only ketones with CH <sub>3</sub> CO— unit give yellow ppt of CHI <sub>3</sub> when heated with aq. alkaline I <sub>2</sub> .







**RAFFLES INSTITUTION**  
**Year 6 H2 CHEMISTRY 2016**  
**Inorganic Chemistry Revision**

<b>1. Summary</b>	
1.1 Periodicity: .....	2
1.2 Group II: .....	4
1.3 Group VII: .....	5
1.4 Transition Elements: .....	7

# 1.1 Periodicity Summary

## Periodicity: Trends Across Period 3

## C. Chlorides (NaCl, MgCl<sub>2</sub>, AlCl<sub>3</sub>, SiCl<sub>4</sub>, PCl<sub>3</sub>/PCl<sub>5</sub>, SCl<sub>2</sub>/S<sub>2</sub>Cl<sub>2</sub>)

### A. Elements (Na, Mg, Al, Si, P, S, Cl, Ar)

#### Physical Properties

##### A1. Atomic radius (Graph A1)

↓ across period due to ↑ Z<sub>eff</sub> as number of protons ↑ but shielding ≈ constant.

##### A2. Ionic Radius (Graph A2)

- Cationic radius (r<sup>+</sup>) ↓ from Na<sup>+</sup> to Al<sup>3+</sup> as Na<sup>+</sup>, Mg<sup>2+</sup> & Al<sup>3+</sup> are isoelectronic but nuclear charge ↑ from Na<sup>+</sup> to Al<sup>3+</sup>.
- Anionic radius (r<sup>-</sup>) ↓ from P<sup>3-</sup> to Cl<sup>-</sup> as P<sup>3-</sup>, S<sup>2-</sup> & Cl<sup>-</sup> are isoelectronic but Z ↑ from P to Cl. Also, ↓ in -ve charge ⇒ ↓ in electronic repulsion.
- r<sub>+</sub> < r<sup>-</sup> ∴ period 3 cations has one shell less than anions.

##### A3. Melting point (Graph A3)

- Element changes from metals (Na, Mg, Al) to non-metals with giant cmolecular structure (Si) and simple molecular structure (P<sub>4</sub>, S<sub>8</sub>, Cl<sub>2</sub> and Ar).
- Mp of Na, Mg and Al are high with mp of Na < Mg < Al ∴ ↑ no. of delocalised e<sup>-</sup> so that metallic bond strength ↑.
- Mp of Si very high ∴ melting involves breaking strong Si-Si bond.
- Mp of P<sub>4</sub>, S<sub>8</sub>, Cl<sub>2</sub> and Ar low ∴ weak id-id interactions between molecules.

##### A4. Electrical Conductivity (Graph A4)

- Na, Mg and Al are metals and are good electrical conductors. Electrical conductivity ↑ from Na to Al ∴ ↑ no. of delocalised e<sup>-</sup>.
- Si is a semi-conductor and conducts electricity minimally.
- P, S, Cl<sub>2</sub> and Ar are insulators of electricity.

#### Chemical Properties

##### A5. 1<sup>st</sup> Ionisation Energy (Graph A5)

- 1<sup>st</sup> IE generally ↑ across period due to ↑ Z<sub>eff</sub> as number of protons ↑ but shielding ≈ constant. 2 exceptions:
- 1) 1<sup>st</sup> IE of Mg > Al because e<sup>-</sup> of Al removed from a 3p orbital which is higher in energy than 3s orbital where e<sup>-</sup> of Mg is removed from.
- 2) 1<sup>st</sup> IE of P > S because e<sup>-</sup> of S removed from a paired orbital and is aided by e<sup>-</sup>e<sup>-</sup> repulsion.

##### A6. Rxn with O<sub>2</sub>

Element	Deg of Rxn	Oxide formed
Na	vigorous	Na <sub>2</sub> O
Mg	vigorous	MgO
Al	vigorous at 1 <sup>st</sup> then slow	Al <sub>2</sub> O <sub>3</sub>
Si	slow	SiO <sub>2</sub>
P	vigorous	P <sub>4</sub> O <sub>6</sub> &/or P <sub>4</sub> O <sub>10</sub>
S	slow	SO <sub>2</sub> &/or SO <sub>3</sub>

##### A8. Variation in Oxidation No. of Oxides and Chlorides

- [O] no. of Na, Mg and Al in oxide and chlorides are fixed at +1, +2 and +3 respectively as the metals loses or shares (only for AlCl<sub>3</sub>) their valence e<sup>-</sup>(s) in bonding.
- Si shares its 4 valence e<sup>-</sup> in oxide and chloride formation so its [O] no. is +4.
- P forms P<sub>4</sub>O<sub>6</sub> & PCl<sub>3</sub> with [O] no. of +3, by sharing 3 e<sup>-</sup> to achieve octet configuration. It can also expand its octet by promoting one 3s e<sup>-</sup> to 3d subshell to form P<sub>4</sub>O<sub>10</sub> and PCl<sub>5</sub> with [O] no. +5
- S forms SO<sub>2</sub> with [O] no. +4 by promoting one 3p e<sup>-</sup> to 3d subshell and SO<sub>3</sub> with [O] no. +6 by promoting one 3s and one 3p e<sup>-</sup> to 3d subshell.

##### B1. Types of Oxides (Rxn with acids and bases):

- Na<sub>2</sub>O & MgO: ionic oxides. Basic oxides that react with acids but not bases:  
Na<sub>2</sub>O(s) + 2H<sup>+</sup>(aq) → 2Na<sup>+</sup>(aq) + H<sub>2</sub>O(l)  
MgO(s) + 2H<sup>+</sup>(aq) → Mg<sup>2+</sup>(aq) + H<sub>2</sub>O(l)
- Al<sub>2</sub>O<sub>3</sub>: ionic oxide with high covalent character. Amphoteric oxide it reacts with acids & bases:  
Al<sub>2</sub>O<sub>3</sub>(s) + 6H<sup>+</sup>(aq) → 2Al<sup>3+</sup>(aq) + 3H<sub>2</sub>O(l)  
Al<sub>2</sub>O<sub>3</sub>(s) + 2OH<sup>-</sup>(aq) + 3H<sub>2</sub>O(l) → 2Al(OH)<sub>4</sub><sup>-</sup>(aq)
- SiO<sub>2</sub>: Covalent oxide with giant molecular structure. Acidic oxide that reacts with conc NaOH:  
SiO<sub>2</sub>(s) + 2OH<sup>-</sup>(aq) → SiO<sub>3</sub><sup>2-</sup>(aq) + H<sub>2</sub>O(l)

##### B2. Reaction of Oxides with Water (Graph B2)

- Na<sub>2</sub>O is ionic & soluble. Reacts with H<sub>2</sub>O to form aq NaOH: (pH ≈ 12)  
Na<sub>2</sub>O(s) + H<sub>2</sub>O(l) → 2NaOH(aq)
- MgO: ionic oxide & sparingly soluble (pH ≈ 9):  
MgO(s) + H<sub>2</sub>O(l) ⇌ Mg(OH)<sub>2</sub>(s) ⇌ Mg<sup>2+</sup>(aq) + 2OH<sup>-</sup>(aq)
- Al<sub>2</sub>O<sub>3</sub> & SiO<sub>2</sub>: insoluble in H<sub>2</sub>O ⇒ after adding Al<sub>2</sub>O<sub>3</sub> & SiO<sub>2</sub> to H<sub>2</sub>O, pH of resultant mixture = 7.
- P<sub>4</sub>O<sub>10</sub>, SO<sub>3</sub> and Cl<sub>2</sub>O<sub>7</sub> reacts with water to give oxoacids (pH ≈ 2)  
P<sub>4</sub>O<sub>10</sub>(s) + 6H<sub>2</sub>O(l) → 4H<sub>3</sub>PO<sub>4</sub>(aq)

##### A7. Rxn with Cl<sub>2</sub>

Element	Deg of Rxn	Chloride formed
Na	vigorous	NaCl
Mg	vigorous	MgCl <sub>2</sub>
Al	vigorous	AlCl <sub>3</sub>
Si	slow	SiCl <sub>4</sub>
P	slow	PCl <sub>3</sub> &/or PCl <sub>5</sub>
S	slow	SCl <sub>2</sub> &/or S <sub>2</sub> Cl <sub>2</sub>

##### C1. Types of Chlorides:


- NaCl & MgCl<sub>2</sub>: ionic chlorides. High mp and bp.
- AlCl<sub>3</sub>, SiCl<sub>4</sub>, PCl<sub>3</sub>/PCl<sub>5</sub>, SCl<sub>2</sub>/S<sub>2</sub>Cl<sub>2</sub> are all covalent with simple molecular structures. Mp and bp low due to weak IMF.
- Bp: AlCl<sub>3</sub> (actually Al<sub>2</sub>Cl<sub>6</sub> ∴ of dimerisation, highest no. of e<sup>-</sup>) > PCl<sub>5</sub> (∴ of pd-pd interactions) > SiCl<sub>4</sub> > S<sub>2</sub>Cl<sub>2</sub>

##### C2. Reaction of Chlorides with Water (Graph C2)


- NaCl is ionic & soluble. Dissolves in H<sub>2</sub>O to give NaCl(aq) : (pH = 7)  
NaCl(s) + aq → NaCl(aq)
- MgCl<sub>2</sub> & AlCl<sub>3</sub> dissolve in water to give Mg<sup>2+</sup> & Al<sup>3+</sup>. The aq ions exist as aqua complexes and hydrolyse in water:  
[Mg(H<sub>2</sub>O)<sub>6</sub>]<sup>2+</sup> ⇌ [Mg(H<sub>2</sub>O)<sub>5</sub>(OH)]<sup>+</sup> + H<sup>+</sup> (pH ≈ 6.5)  
[Al(H<sub>2</sub>O)<sub>6</sub>]<sup>3+</sup> ⇌ [Al(H<sub>2</sub>O)<sub>5</sub>(OH)]<sup>2+</sup> + H<sup>+</sup> (pH ≈ 3)
- SiCl<sub>4</sub>, PCl<sub>3</sub> and S<sub>2</sub>Cl<sub>2</sub> hydrolyses in water: (pH ≈ 2 ∴ HCl is strong acid.)  
SiCl<sub>4</sub>(l) + 2H<sub>2</sub>O(l) → SiO<sub>2</sub>(s) + 4HCl(aq)

Refer to Graph B3 and C3 for melting points of oxides and chlorides


A1: Variation of Atomic Radius for Period 3 Elements




A2: Variation of Ionic Radius for Period 3 Elements




A3: Variation of Melting Points for Period 3 Elements




A4: Variation of Electrical Conductivity for Period 3 Elements




B2: Variation of pH of resultant mixture when Period 3 Oxide is added to water




B3: Variation of Melting Points for Period 3 Oxides




A4: Variation of 1<sup>st</sup> Ionisation Energy for Period 3 Elements



C2: Variation of pH of resultant mixture when Period 3 Chloride is added to water



C3: Variation of Melting Points for Period 3 Chlorides



# 1.2 Group II Summary

## A. Elements:

Be, Mg, Ca, Sr, Ba

### A1. Reactivity as Reducing Agent:

- Group II metals lose  $e^-$  in their rxn:  
 $M \rightarrow M^{2+} + 2e^-$ , ie they become oxidised and hence act as reducing agents (RA)
- Strength of a RA depends on its ease to be oxidised.
- Greater ease of oxidation  $\Rightarrow$  greater strength as RA.
- Since [O] involves loss of  $e^-$ , strength of RA  $\uparrow$  when ease of loss of  $e^- \uparrow$ .
- Ease of loss of  $e^- \uparrow$  down Gp II as valence  $e^-$  is further from nucleus and shielded by more filled inner quantum shells.
- Hence reactivity as RA  $\uparrow$  down Gp II.

### A2. Reaction with Water:

Element	Deg of Rxn	Equations
Mg	Very slow rxn with cold water.	-
Ca	Readily	$M(s) + 2H_2O(l) \rightarrow$
Sr	Vigorously	$M(OH)_2$
Ba	Very vigorously	$(aq) + H_2(g)$

### A3. Reaction with Air:

Element	Deg of Rxn	Equations
Mg	Ignites above mp	
Ca	Ignites on heating	$2M(s) + O_2(g) \rightarrow$
Sr	Ignites	$2MO(s)$
Ba	spontaneously	

## Group II Chemistry

### B. Type of Gp II oxides

- BeO is amphoteric and reacts with both acids and bases:  
 $BeO + 2H^+ \rightarrow Be^{2+} + H_2O$   
 $BeO + 2OH^- \rightarrow Be(OH)_4^{2-}$
- All other gp II oxides are basic and reacts with acids but not bases.

### C. Thermal Stability of Group II Cpds containing oxoanions

- Thermal stability  $\uparrow$  down Gp II.
- Cation size  $\uparrow$  down Gp II resulting in  $\downarrow$  in charge density and hence polarising power.
- Anion electron cloud is polarised to a smaller extent by cation resulting in  $\downarrow$  weakening of covalent bond in anion.
- Hence a higher temperature is needed for decomposition of oxoanion to oxide.
- Decomposition of Nitrate on Heating:  
 $M(NO_3)_2(s) \rightarrow MO(s) + 2NO_2(g) + \frac{1}{2}O_2(g)$
- Decomposition of Carbonate on Heating:  
 $MCO_3(s) \rightarrow MO(s) + CO_2(g)$

### D. Uses of Gp II Compounds

- MgO used as refractory material to line furnaces due to its high melting point.
- Lime (CaO) used to neutralize acidic soil.
- NB: Lime/Quicklime = CaO; Slaked lime =  $Ca(OH)_2(s)$ ; Limewater =  $Ca(OH)_2(aq)$ ; Limestone =  $CaCO_3$ .

## Solubility of common ionic compounds:

- All ammonium, potassium and sodium compounds are soluble.
- All nitrates are soluble.
- All halides ( $X^- = Cl^-, Br^-$  or  $I^-$ ) are soluble except AgX and  $PbX_2$ .
- All sulphates are soluble except  $PbSO_4$  and  $BaSO_4$ .
- All carbonates are insoluble except carbonates of  $NH_4^+$ ,  $K^+$  and  $Na^+$ .
- All hydroxides are insoluble except Gp I hydroxides (eg NaOH and KOH) and  $Ca(OH)_2$ ,  $Sr(OH)_2$  and  $Ba(OH)_2$ . Metal oxides have similar solubility as metal hydroxides.

## Reactions of Common Cations with aq NaOH:

- No ppt, no  $NH_3$  on warming  $\Rightarrow H^+$ ,  $Na^+$ ,  $K^+$
- No ppt,  $NH_3$  on warming  $\Rightarrow NH_4^+$
- White ppt insoluble in xs aq NaOH  $\Rightarrow$  Group II cations:  $Mg^{2+}$ ,  $Ca^{2+}$ ,  $Ba^{2+}$
- White ppt soluble in xs aq NaOH  $\Rightarrow Pb^{2+}$ ,  $Al^{3+}$ ,  $Zn^{2+}$
- White ppt insoluble in xs aq NaOH, turns brown on standing  $\Rightarrow Mn^{2+}$
- Greyish green ppt soluble in xs aq NaOH  $\Rightarrow Cr^{3+}$
- Dirty green ppt, insoluble in xs aq NaOH  $\Rightarrow Fe^{2+}$
- Reddish brown ppt, insoluble in xs aq NaOH  $\Rightarrow Fe^{3+}$
- Blue ppt, insoluble in xs aq NaOH  $\Rightarrow Cu^{2+}$

## Reactions of Common Cations with aq $NH_3$ :

- No ppt  $\Rightarrow H^+$ ,  $Na^+$ ,  $K^+$ ,  $Ca^{2+}$ ,  $Ba^{2+}$
- White ppt insoluble in xs aq  $NH_3 \Rightarrow Mg^{2+}$ ,  $Pb^{2+}$ ,  $Al^{3+}$
- White ppt soluble in xs aq  $NH_3 \Rightarrow Zn^{2+}$  [due to formation of  $Zn(NH_3)_4^{2+}(aq)$ ]
- White ppt insoluble in xs aq  $NH_3$ , turns brown on standing  $\Rightarrow Mn^{2+}$
- Greyish green ppt, insoluble in xs aq  $NH_3 \Rightarrow Cr^{3+}$
- Dirty green ppt, insoluble in xs aq  $NH_3 \Rightarrow Fe^{2+}$
- Reddish brown ppt, insoluble in xs aq  $NH_3 \Rightarrow Fe^{3+}$

## Test for Anions:

- $CO_3^{2-}$ : reacts with  $H^+$  to give  $CO_2$ .
- $SO_3^{2-}$ : reacts with  $H^+$  to give  $SO_2$ .
- $NO_3^-$ : reacts with Al foil and aq NaOH on heating to give  $NH_3$ .
- $SO_4^{2-}$ : reacts with aq  $Ba(NO_3)_2$  to give white ppt, insoluble in dil  $HNO_3$ .
- $CrO_4^{2-}$ : turns from yellow to orange on addition of  $H^+$ .
- $Cl^-$ : reacts with aq  $AgNO_3$  to give white ppt, insoluble in dil  $HNO_3$ .
- $Br^-$ : reacts with aq  $AgNO_3$  to give cream ppt, insoluble in dil  $HNO_3$ .
- $I^-$ : reacts with aq  $AgNO_3$  to give yellow ppt, insoluble in dil  $HNO_3$ .

## Test for Gases:

- $H_2$ : extinguishes lighted splinter with a 'pop' sound.
- $O_2$ : relights a glowing splinter
- $CO_2$ : forms a white ppt with aq  $Ca(OH)_2$
- $SO_2$ : decolourises aq  $KMnO_4$
- $Cl_2$ : turns moist blue litmus red then bleach it.
- $NH_3$ : turns moist red litmus blue.

## 1.3 GROUP VII SUMMARY

**Overview:** Key reactivity for halogens is redox reactions.

- (1) Oxidizing power trend
- (2) Reaction with hydrogen
- (3) Displacement of halide by more reactive halogen

(4) Reaction with water

(oxidation of water, disproportionation)

(5) Reaction with thiosulfate

(oxidize S O.S +2 → +6 or +2.5)

(6) Disproportionation in alkalis

(O.S → +1, -1, or +5, -1)

Aside from F<sub>2</sub>, typically Cl<sub>2</sub> is most reactive and most oxidizing.




It is best to have the skills to derive and balance the equations quickly, rather than memorizing them by rote

### (1) Oxidizing power

A substance that is readily reduced would oxidize another substance in a redox reaction. => top of the group: more +ve E, higher oxidizing power.

Best oxidizing agent [OA]

↑



↑

Weak Ox. agent

Good reducing agent

(since backward rxn takes place easily)

### (2) Reaction with H<sub>2</sub>

All react to give HX.

F<sub>2</sub> violent explosion. No heat or light required.

Cl<sub>2</sub>. Violent explosion. UV light or heat to start reaction.

Br<sub>2</sub>. Slow reaction. Heat and catalyst required.

I<sub>2</sub>. Slow, incomplete/reversible reaction.

### (3) Displacement of halide by more reactive halogen

Cl<sub>2</sub> displaces Br<sup>-</sup>, I<sup>-</sup>. Br<sub>2</sub> displaces I<sup>-</sup>.

Eg:  $Cl_2 + 2Br^- \rightarrow 2Cl^- + Br_2$


### (4) Reaction with water

F<sub>2</sub> oxidizes H<sub>2</sub>O to give O<sub>2</sub> and HF

Cl<sub>2</sub> disproportionates to +1 and -1 O.S.

Br<sub>2</sub> does not react.

I<sub>2</sub> dissolves in I<sup>-</sup>(aq) by forming I<sub>3</sub><sup>-</sup>(aq)



### How to remember the equations?

- For redox reactions, you need to know the oxidation states => formula of starting materials and products.

- If O.S is known, the identity of the halide/halate can be determined. O.S of X increases by +2 for every O atom added. i.e.

-1	+1	+3	+5	+7
X <sup>-</sup>	XO <sup>-</sup>	XO <sub>2</sub> <sup>-</sup>	XO <sub>3</sub> <sup>-</sup>	XO <sub>4</sub> <sup>-</sup>

- The balanced equations can easily be derived.

\*Note. Fast redox eqn balancing method

### Balancing Redox equations

\*Note. The best way to remember the reactions is to note: (1) type of reaction., (2) for redox, note the Oxidation state(O.S) reached.

(1) use ΔO.S for each atom oxidized or reduced to determine no. of e<sup>-</sup> transferred.

(2) use LCM(lowest common multiple) of the no. of e<sup>-</sup> transferred to find stoichiometric ratio of [R] and [O] agents.


(3) balance LHS and RHS charge with charged particles available (usually H<sup>+</sup> or OH<sup>-</sup>)

(4) balance O and H atoms with H<sub>2</sub>O last.


See next section for example with S<sub>2</sub>O<sub>3</sub><sup>2-</sup>

### (5) Reaction with thiosulfate

Cl<sub>2</sub> and Br<sub>2</sub> strong OA, will oxidize S from +2 to +6




I<sub>2</sub> weaker [OA], oxidize S from +2 to +2.5



\*Standard and in databook.


Example of how to balance equations:




(0) (+2) (-1) (+6)

Cl(0 → -1) 1 eln gain; S(+2 → +6) 4 eln lost.


So LCM is 4, ratio of Cl<sub>2</sub>:S is 4:1 Thus



(3) Balance charge with H<sup>+</sup>. excess 10 -ve charge on RHS so:




(4) Balance O and H with H<sub>2</sub>O. so:



### (6) Disproportionation in alkali


Cl<sub>2</sub>, good OA, but difficult to oxidize. so at rtp disproportionates in alkaline solutions.

Cold or rtp, Cl<sub>2</sub> 0 → +1 and -1



On warming. Further disproportionation

+1 → -1 and +5




Overall/or Cl<sub>2</sub> in hot OH<sup>-</sup>: 0 → -1 and +5



Br<sub>2</sub>, I<sub>2</sub> more easily oxidized to higher O.S

0 → +5 and -1




T > 15 °C for Br<sub>2</sub>.

### How to remember the equations?

Same redox method used in previous section can be used for any reaction here.


Example of how to balance equations:




(0) (-1) (+1)

Cl(0 → -1) 1 eln gain; Cl(0 → +1) 1 eln lost.


So LCM is 1, ratio of Cl<sup>-</sup>:ClO<sup>-</sup> is 1:1 Thus



(3) Balance charge with OH<sup>-</sup>. excess 2 -ve charge on RHS so:




(4) Balance O and H with H<sub>2</sub>O. so:




## X<sub>2</sub> in hot OH<sup>-</sup>

Example of how to balance equations:




So LCM is 5, ratio of  $\text{Cl}^- : \text{ClO}_3^-$  is 5:1 Thus




Balance Cl atoms  $3\text{Cl}_2 \rightarrow 5\text{Cl}^- + \text{ClO}_3^-$

(3) Balance charge with  $\text{OH}^-$ . excess 6 -ve charge on RHS so:



(4) Balance O and H with  $\text{H}_2\text{O}$ . so:



If counter-ion is to be included, simply add to balance off against all ions. For eg. if NaOH was used, add  $\text{Na}^+$  to balance of all the anions.

Thus:




## (7) Physical Properties of halogens

- Colour intensifies down the group.

### Boiling point

- b.p. increases down the group as size of electron cloud increases, van der Waals' forces between  $\text{X}_2$  molecules increase.



### X-X Covalent bond strength.

- F-F bond is unusually weak. The F-F bond is very short and the atoms are close together. The lone pair electrons on both atoms repel and push the atoms apart, weakening the bond.

- The X-X bond down the group weakens, since the atoms become larger down the group and orbital overlap is less effective. Thus Cl-Cl bond is strongest amongst  $\text{X}_2$  molecules.

X-X covalent bond strength:  $\text{F-F} < \text{Cl-Cl} > \text{Br-Br} > \text{I-I}$

## Properties of halides

(1) hydrogen halides: physical properties, acidity

(2) halide salts,

- reaction with conc.  $\text{H}_2\text{SO}_4$


- precipitation reactions ( $\text{Ag}^+$ )

## (1) Hydrogen halides

### Boiling point

- b.p is especially high for H-F, since there are strong hydrogen bonds between HF molecules.

- b.p is lowest for  $\text{H-Cl} < \text{H-Br} < \text{H-I}$ . Since the size of electron cloud increases down the group, the strength of the van der Waals' forces increase down the group for HX.



### H-X covalent bond strength

The H-X covalent bond weakens down gp VII. As the X atom becomes larger, orbital overlap with H atom becomes poorer.

Cov bond strength:  $\text{H-F} > \text{H-Cl} > \text{H-Br} > \text{H-I}$

### Stability

- H-Br decomposes slightly with heat.

- H-I is least stable and decomposes easily with heat

### Acidity

- H-F bond is strongest and so HF is a weaker acid, ionizing less readily.

- HCl, HBr, HI are all strong acids and ionize completely in water. However relatively, acid strength of  $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$

## (2) Halide salts

### Reaction with c. $\text{H}_2\text{SO}_4$

$\text{H}_2\text{SO}_4$  has 2 roles

- Acid  $\Rightarrow$  acid base reaction possible

- S atom at +6 O.S. Thus it can be reduced, and has limited oxidizing power.

Thus

(1)

With any halide salt eg. NaX, protonation to HX occurs first:





White fumes evolved.

(2)




$\text{HCl/Cl}^-$  has a poor reducing power and no further redox reactions observed.

$\text{HBr/Br}^-$  has moderate reducing power,



Reddish brown liquid, fumes evolved.



$\text{HI/I}^-$  has good reducing power, and will reduce S atom in  $\text{H}_2\text{SO}_4$  ( $\text{O.S } +6 \rightarrow +4, 0, -2$ ). Thus:



### How to remember the equations?


Same redox method used in previous section can be used for any reaction here.

Example, HI with  $\text{H}_2\text{SO}_4$  to reduce S ( $\text{OS } +6 \rightarrow -2$ )



$\text{I}(-1 \rightarrow 0) \text{ 1 eln lost; S}(+6 \rightarrow -2) \text{ 8 eln gain.}$


So LCM is 8, ratio of I:S is 8:1 Thus



(make sure all the I and S atoms are balanced)


(3) Balance charge: not necessary since all neutral

(4) Balance O and H with  $\text{H}_2\text{O}$ . so:



### Precipitation of silver halides, AgX



All halides form precipitates with  $\text{Ag}^+$ :



$\text{AgCl}$  – white ppt,  $\text{AgBr}$  – cream ppt,  $\text{AgI}$  – yellow ppt

$\text{AgI}$  has the smallest  $K_{sp}$  and lowest solubility.

Excess  $\text{NH}_3(\text{aq})$  can dissolve  $\text{AgCl}$  by forming silver diammine



$[\text{Ag}^+(\text{aq})]$  drops and IP falls below  $K_{sp}$  of  $\text{AgCl}$  hence,  $\text{AgCl}$  ppt dissolves.

## 1.4 Transition Elements Summary

### First Row Transition Elements

A transition element is defined as an element that possesses an incomplete d subshell in at least one of its stable ions. Scandium is not a transition metal as its only common ion  $\text{Sc}^{3+}$  does not possess d electron. Zinc is also not a transition metal as it contains a full 3d subshell in its only oxidation state +2. Note: Cr and Cu has one 4s electron only as it is more energetically favourable for them to have a half-filled or fully filled 3d subshell.

Element	(Sc)	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	(Zn)
Electronic configuration	$[\text{Ar}]3d^1 4s^2$	$[\text{Ar}]3d^2 4s^2$	$[\text{Ar}]3d^3 4s^2$	$[\text{Ar}]3d^5 4s^1$	$[\text{Ar}]3d^5 4s^2$	$[\text{Ar}]3d^6 4s^2$	$[\text{Ar}]3d^7 4s^2$	$[\text{Ar}]3d^8 4s^2$	$[\text{Ar}]3d^{10} 4s^1$	$[\text{Ar}]3d^{10} 4s^2$

Essentially for TEs, the special properties arise from the presence of d-electrons and d orbitals: eg catalytic property, colour.

### Physical Trends

1<sup>st</sup> 3 TMs have smaller atomic radii than s-block metals: S-block metals use inner shell s and p electrons for shielding while s, p and d electrons are used for shielding in TMs. d-orbitals are more diffuse and the d electrons provide poor shielding effect; thus TMs experience a higher effective nuclear charge.

Later TMs have almost constant atomic radii and 1<sup>st</sup> I.E.: Across the d-block, nuclear charge increases but shielding effect increases as additional electrons are added to the penultimate (inner) 3d subshell. The increase in shielding effect offsets the increase in nuclear charge and effective nuclear charge increases gradually. Hence the atomic radii and first I.E. are almost constant.

Note there is no contradiction between the first two points. For the latter TMs, the 4s electrons are repelled out, as well as the d-subshell having a higher electron density; thus shielding effect from d electrons is insignificant in earlier TMs while significant for latter TMs.

TMs have higher melting points and boiling points as well as higher electrical conductivity: The 4s and 3d electrons are available for delocalization, hence there are more delocalized electrons and cations have higher charge. Metallic bonding is stronger due to these and smaller cationic radii compared to s-block metals. There are also more delocalized electrons to conduct electricity.

### Variable Oxidation State

TMs have variable oxidation states (OS) as the 3d and 4s electrons are very similar in energy. When TMs lose electrons, they lose the 4s electrons first and the +2 OS is common. The  $\text{M}^{4+}$  ion is rare and in higher oxidation states, the element is usually not found in the free metal ion but covalently bonded or as oxyanions, e.g.  $\text{MnO}_4^-$ .

### Catalytic Properties

Catalysts: provide alternate reaction path with lower  $E_a$ . Itself not consumed in net reaction. 2 types:

Homogeneous: Same phase (eg in aq solution) involves rapid conversion from one oxidation state to another oxidation state. Eg.  $\text{Fe}^{2+}$  in the reaction between  $\text{I}^-$  and  $\text{S}_2\text{O}_8^{2-}$ .

The metal ion acts as an electron carrier. Higher OS cation oxidizes one reagent, while lower OS cation reduces another.

-uncatalysed reaction: slow because of high  $E_a$ . For eg  $\text{I}^-$  and  $\text{S}_2\text{O}_8^{2-}$  both -ve charged and repel.

-catalysed reaction: multiple steps, new reaction path.

Eg:  $\text{Fe}^{3+}$  oxidizes  $\text{I}^-$  to give  $\text{Fe}^{2+}$  and  $\text{I}_2$ .

$\text{Fe}^{2+}$  then reduced  $\text{S}_2\text{O}_8^{2-}$  to  $\text{SO}_4^{2-}$ . Each step has lower  $E_a$ , since oppositely charged ions which attract are involved, thus the reaction is faster.  $E^\circ_{\text{cell}}$  for each step can be calculated to show feasibility.

Heterogeneous: Depends on the availability of vacant low-lying orbitals. and availability of d electrons for bonding with reactants. E.g. Fe in Haber process, Ni in hydrogenation of alkenes,  $\text{MnO}_2$  in decomposition of  $\text{H}_2\text{O}_2$ ,  $\text{V}_2\text{O}_5$  in Contact Process.

-diffusion, adsorption on surface (bond breaking within reagent molecules, effective orientation of reactants), -reaction (new bonds form to give products), -diffusion(product molecules leave)

### Formation of Complex Ions

TM ions have high charge densities and can attract ligands, which are neutral molecules or anions containing lone pairs. The ligands form dative covalent bonds with the central metal ion to form complex ions.

Coordination Number	Examples	Shape
6	$[\text{Fe}(\text{CN})_6]^{3-}$ $[\text{Cu}(\text{edta})]^{2-}$	Octahedral
4	$[\text{CuCl}_4]^{2-}$ $[\text{Cu}(\text{NH}_3)_4]^{2+}$	Square planar or tetrahedral
2	$[\text{Ag}(\text{NH}_3)_2]^+$ , $[\text{CuCl}_2]^-$	Linear

Key concepts:

1) Ligand exchange. Takes place when more stable complexes can be formed. Expect to see colour change, etc

2)  $E^\circ$  will be different. For eg  
 $\text{Fe}^{3+}(\text{aq}) (E^\circ = +0.77 \text{ V})$  vs  $[\text{Fe}(\text{CN})_6]^{3-}(\text{aq}) (E^\circ = +0.36 \text{ V})$   
 -ve charge complex repels electrons and less easily reduced


$\text{Fe}^{3+}(\text{aq}) (E^\circ = +0.77 \text{ V})$  vs  $\text{Fe}(\text{OH})_3(\text{s}) (E^\circ = -0.56 \text{ V})$   
 Stronger ligand (the  $\text{OH}^-$ ) increases electron density on metal centre, decreased ease of reduction. Also the Fe(III) centre is stabilized by electron donated from ligand.

3) Formation of complex ions can "extract" the metal ion from solution and thus enable the ppt of the ion to dissolve. Eg:

-excess conc  $\text{Cl}^-$  dissolves  $\text{PbCl}_2$ , forms  $[\text{PbCl}_4]^{2-}$

- addition of aq.  $\text{NH}_3$  dissolves  $\text{AgCl}$  ppt., forms  $[\text{Ag}(\text{NH}_3)_2]^+$

- excess  $\text{NH}_3$  dissolves  $\text{Cu}(\text{OH})_2$ . Net reaction:



## Coloured Complexes

In the gaseous or free ion, the five d orbitals are degenerate (of equal energy). In the presence of ligands, the d orbitals are split into two distinct levels, creating an energy gap. The energy difference between the levels corresponds to a particular wavelength in the visible region of the spectrum. When light falls on the complex, light with wavelength and energy corresponding to the energy gap are absorbed and the d electrons are promoted from the lower energy d orbitals to the higher higher energy d orbitals (d-d transition). The complex exhibits the colour complementary to that of the absorbed wavelengths.

The splitting of the d orbitals depends on the nature of the TM, its oxidation state, the shape of the complex and the nature of the ligands, hence different complexes have different colours. If the d orbitals are completely empty ( $\text{Sc}^{3+}$ ) or completely full ( $\text{Cu}^+$  or  $\text{Zn}^{2+}$ ), no transitions within the d levels can take place and the complexes are colourless.

Vanadium	
$\text{VO}_2^+ / [\text{VO}_2(\text{H}_2\text{O})_4]^+$	yellow
$\text{VO}^{2+} / [\text{VO}(\text{H}_2\text{O})_5]^{2+}$	blue
$\text{V}^{3+} / [\text{V}(\text{H}_2\text{O})_6]^{3+}$	green
$\text{V}^{2+} / [\text{V}(\text{H}_2\text{O})_6]^{2+}$	violet
$\text{VF}_5$	white
$\text{VF}_4$	lime green
$\text{VF}_3$	yellow green
$\text{VF}_2$	blue
$\text{VCl}_4$	red-brown
$\text{VCl}_3$	red-violet
$\text{VCl}_2$	pale green
Chromium	
$\text{Cr}^{2+} / [\text{Cr}(\text{H}_2\text{O})_6]^{2+}$	blue/sky-blue
$\text{Cr}^{3+} / [\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ (aq)	green
$[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ solid salt	violet
$\text{CrO}_4^{2-}$	yellow
$\text{Cr}_2\text{O}_7^{2-}$	orange
$\text{Cr}(\text{OH})_3 / [\text{Cr}(\text{OH})_3(\text{H}_2\text{O})_3]$	green ppt
$[\text{Cr}(\text{OH})_6]^{3-}$	deep green
Manganese	
$\text{Mn}^{2+} / [\text{Mn}(\text{H}_2\text{O})_6]^{2+}$	pale pink
$\text{Mn}(\text{OH})_2$	off-white ppt/buff ppt
$\text{Mn}_2\text{O}_3 \cdot x\text{H}_2\text{O}$	brown ppt
$\text{MnCl}_4^{2-}$	green-yellow
$\text{MnO}_2$	black solid
$\text{MnO}_4^-$	deep purple
$[\text{Mn}(\text{CN})_6]^{3-}$	dark red
Iron	
$[\text{Fe}(\text{CN})_6]^{3-}$	orange-brown
$[\text{Fe}(\text{CN})_6]^{4-}$	yellow
$\text{Fe}(\text{OH})_3 / [\text{Fe}(\text{OH})_3(\text{H}_2\text{O})_3]$	reddish brown
$\text{Fe}(\text{OH})_2 / [\text{Fe}(\text{OH})_2(\text{H}_2\text{O})_4]$	dirty green
$\text{Fe}_2\text{O}_3 \cdot x\text{H}_2\text{O}$	reddish brown
$\text{FeCO}_3$	white
$\text{Fe}^{3+} / [\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ (aq)	yellow
$[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ (solid salt)	lilac
$[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$	pale green
$[\text{Fe}(\text{SCN})(\text{H}_2\text{O})_5]^{2+}$	blood red
$[\text{Fe}(\text{CN})_6]^{4-} + \text{Fe}^{2+}$	Prussian blue ppt
$[\text{Fe}(\text{CN})_6]^{3-} + \text{Fe}^{3+}$	Turnbull's blue ppt
$\text{Fe}_4[\text{Fe}(\text{CN})_6]_3 \cdot 15\text{H}_2\text{O}$	dark blue ppt (same as above)

Cobalt	
$[\text{Co}(\text{NH}_3)_6]^{3+}$	golden brown
$[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{O})]^{3+}$	bright red
$[\text{Co}(\text{NH}_3)_5\text{Cl}]^{2+}$	purple
$[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$ cis	deep violet
$[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$ trans	green
$[\text{CoCl}_4]^{2-}$	blue
$[\text{Co}(\text{H}_2\text{O})_6]^{2+}$	pink
$\text{Co}(\text{OH})_2$	blue ppt
$[\text{Co}(\text{NH}_3)_6]^{2+}$	pale brown
Nickel	
$[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$	green
$[\text{Ni}(\text{NH}_3)_6]^{2+}$	blue
$[\text{Ni}(\text{CN})_4]^{2-}$ sq planar	yellow
$[\text{NiCl}_4]^{2-}$	blue
$\text{Ni}(\text{OH})_2 / [\text{Ni}(\text{OH})_2(\text{H}_2\text{O})_4]$	pale green ppt
$\text{Ni}(\text{DMG})$	red ppt
DMG = dimethylglyoxime	
Copper	
$\text{Cu}(\text{I})$ compounds (except below)	white solids
$\text{Cu}_2\text{O}$	red ppt
$\text{Cu}_2\text{S}$	black solid
$\text{CuI}$	cream ppt
$\text{CuCN}$	white ppt
$[\text{Cu}(\text{CN})_4]^{3-}$	colorless
$[\text{CuCl}_2]$	colorless
$\text{CuO}$	black solid
$\text{CuCO}_3$	green solid
$\text{CuCl}_2$ (anhydrous)	dark brown
$\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$	blue-yellow
$\text{Cu}^{2+} / [\text{Cu}(\text{H}_2\text{O})_6]^{2+}$	pale blue
$[\text{Cu}(\text{NH}_3)_4]^{2+} / [\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$	deep blue
$\text{Cu}(\text{OH})_2 / [\text{Cu}(\text{OH})_2(\text{H}_2\text{O})_4]$	blue ppt
$[\text{CuCl}_4]^{2-}$	yellow
$[\text{Cu}(\text{edta})]^{2-}$	pale blue / light blue
Silver	
$\text{Ag}(\text{I})$ charged complexes	colorless usually
$[\text{Ag}(\text{H}_2\text{O})_2]^+$ , $[\text{Ag}(\text{NH}_3)_2]^+$ , $[\text{Ag}(\text{S}_2\text{O}_3)_2]^{3-}$ , $[\text{Ag}(\text{CN})_2]^-$	colorless
$\text{AgCl}$	white ppt
$\text{AgBr}$	cream ppt
$\text{AgI}$	yellow
$\text{Ag}_2\text{S}$	black ppt
Metal Ion Centre	Usual Co-ord. No.
$\text{Cu}^+$ , $\text{Ag}^+$ ( $d^{10}$ )	2 (linear)
$\text{Cu}^{2+}$ ( $d^9$ )	4 (tetrahedral or sq. planar)
$\text{Ni}^{2+}$ , $\text{Pt}^{2+}$ ( $d^8$ )	4 (sq. planar)
Others	6 (octahedral)